

STATUS OF  
**LEOPARDS**  
IN INDIA



**2022**



Status of  
**Leopards**  
in India

---

2022



**Citation:** Qamar Qureshi, Yadvendradev V. Jhala, Satya P. Yadav, Virendra R. Tiwari, Rajendra Garawad and Amit Mallick (eds) 2024. Status of Leopards in India, 2022. National Tiger Conservation Authority, Government of India, New Delhi, and Wildlife Institute of India, Dehradun.

**ISBN No:** 81-85496-86-2

**Cover photograph :**  
Gurinderjit Singh







मंत्री  
पर्यावरण, वन एवं जलवायु परिवर्तन  
और  
श्रम एवं रोजगार  
भारत सरकार



सत्यमेव जयते  
भूपेन्द्र यादव  
BHUPENDER YADAV

MINISTER  
ENVIRONMENT, FOREST AND CLIMATE CHANGE  
AND  
LABOUR & EMPLOYMENT  
GOVERNMENT OF INDIA



संदेश

वन्यजीव संरक्षण के क्षेत्र में, प्रोजेक्ट टाइगर लंबे समय से वन पारिस्थितिकी तंत्र की सुरक्षा का पर्याय बन गया है। यह इस बात का उदाहरण है कि कैसे बाघों के अलावा अन्य प्रजातियों के संरक्षण पर भी ध्यान दिया जाता है। तेंदुओं की स्थिति पर यह रिपोर्ट इस पहल में शामिल व्यापक संरक्षण प्रयासों के प्रमाण के रूप में खड़ी है।

मानव-प्रधान परिदृश्यों में तेंदुए, पर्यावरणीय चुनौतियों के सामने वन्यजीवों के अनुकूलनशीलता का प्रतीक हैं। तेंदुओं के संरक्षण के प्रयास संरक्षित क्षेत्रों की सीमा से परे प्रजातियों की सुरक्षा की प्रतिबद्धता को स्पष्ट रूप से उजागर करते हैं। विशेष सराहना हमारे समर्पित वन विभाग की है जिनके अटूट प्रयास इस उद्देश्य में महत्वपूर्ण योगदान देते हैं।

प्रोजेक्ट टाइगर, बाघों से परे अपना ध्यान केंद्रित करके, पारिस्थितिक तंत्र के अंतर्संबंध और विविध प्रजातियों के संरक्षण को रेखांकित करता है। यह रिपोर्ट भारत सरकार के सहयोगात्मक प्रयासों, वैज्ञानिक कठोरता और संरक्षण की दिशा में गंभीर प्रयासों का प्रदर्शन है। मैं मंत्रालय, राष्ट्रीय बाघ संरक्षण प्राधिकरण के नेतृत्व, राज्य वन विभाग के कर्मचारियों और भारतीय वन्यजीव संस्थान के वैज्ञानिकों सहित इस प्रयास में शामिल सभी लोगों को बधाई देता हूँ।

भारत के यशस्वी प्रधान मंत्री, श्री नरेन्द्र मोदी जी के नेतृत्व एवं मार्गदर्शन में, हम न केवल वन्यजीव का संरक्षण कर रहे हैं, बल्कि हम उस लोकाचार का उदाहरण दे रहे हैं जो हम सभी को एक साथ बांधता है; एक पृथ्वी, एक परिवार और एक भविष्य।

तारीख : 19 .02.2024

  
(भूपेन्द्र यादव)



अश्विनी कुमार चौबे  
Ashwini Kumar Choubey



राज्य मंत्री  
पर्यावरण, वन एवं जलवायु परिवर्तन  
उपभोक्ता मामले, खाद्य और सार्वजनिक वितरण  
भारत सरकार  
MINISTER OF STATE  
ENVIRONMENT, FOREST AND CLIMATE CHANGE  
CONSUMER AFFAIRS, FOOD & PUBLIC DISTRIBUTION  
GOVERNMENT OF INDIA

जब हम "वसुधैव कुटुंबकम" की बात करते हैं, सर्वे भवन्तु सुखिनः सर्वे संतु निरामयाः कहते हैं तो उस परिधि में मात्र मनुष्य ही नहीं, इस वसुधा के सभी प्राणी आते हैं। हमारे यशस्वी प्रधानमंत्री श्री नरेंद्र मोदी जी के दूरदर्शी नेतृत्व में सभी प्राणियों के महत्व को रेखांकित किया है। तेंदुओं पर आधारित यह स्थिति रिपोर्ट (आबादी का आकलन) इस दर्शन का एक ज्वलंत उदाहरण है। भारत के वन्यजीवों के बीच, कोई अन्य वन्यप्राणी तेंदुए से बेहतर मानव-वन्यजीव सह-अस्तित्व का उदाहरण नहीं देता है।

भारत में तेंदुआ की आबादी निरन्तर बढ़ रही है, जिसमें इसकी बड़ी सफलता का श्रेय बाघ संरक्षण परियोजना के अन्तर्गत किए गए सुरक्षा एवं संरक्षण उपायों को जाता है।

भारतीय समुदायों के अतिरिक्त पृथ्वी पर कहीं और कोई भी अन्य समुदाय, वन्य जीवों के प्रति इतनी सहनशीलता नहीं दिखाते हैं। जैव विविधता में गिरावट से जूझ रही दुनिया में, भारत आशा की किरण के रूप में खड़ा है, जो यह दर्शाता है कि एकजुट प्रयासों से वन्यजीवों का संरक्षण हो सकता है।

मैं दुनिया को पर्यावरण संरक्षण एवं वन्यजीव सुरक्षा का सच्चा सार दिखाने के लिए भारत की जनता और हमारे समर्पित वन कर्मियों को हार्दिक बधाई देता हूँ। इस संरक्षण के प्रयास को वास्तविकता बनाने में उनकी महत्वपूर्ण भूमिका के लिए राष्ट्रीय बाघ संरक्षण प्राधिकरण के अधिकारियों, राज्य वन विभाग के कर्मचारियों और भारतीय वन्यजीव संस्थान के वैज्ञानिकों को हार्दिक साधुवाद देता हूँ।

मैं आशा करता हूँ कि हम सब साथ मिलकर आने वाली पीढ़ियों के लिए अपने साझा पर्यावरण एवं वन्यजीवों को संरक्षित एवं संवर्धित करने में सामूहिक उत्तरदायित्व का पालन करेंगे।

(अश्विनी कुमार चौबे)

कार्यालय : 5वां तल, आकाश विंग, इंदिरा पर्यावरण भवन, जोर बाग रोड, नई दिल्ली-110003, दूरभाष : 011-20819418, 011-20819421, फैक्स : 011-20819207, ई-मेल : mos.akc@gov.in  
Office : 5th Floor, Aakash Wing, Indira Paryavaran Bhawan, Jor Bagh Road, New Delhi-110003, Tel.: 011-20819418, 011-20819421, Fax: 011-20819207, E-mail : mos.akc@gov.in

कार्यालय : कमरा नं.173, कृषि भवन, नई दिल्ली-110001, दूरभाष : 011-23380630, फैक्स : 011-23380632

Office : Room No. 173, Krishi Bhawan, New Delhi-110001, Tel. : 011-23380630, Fax : 011-23380632

निवास : 30, डॉ. एपीजे अब्दुल कलाम रोड, नई दिल्ली-110003, दूरभाष : 011-23794971, 23017049

Residence : 30, Dr. APJ Kalam Road, New Delhi-110003, Tel.: 011-23794971, 23017049





# Contents

Executive Summary .....	<b>I-VIII</b>
I. Status of Leopards in India .....	<b>01</b>
II. Shivalik Hills and Gangetic Plains .....	<b>17</b>
III. Central India and Eastern Ghats .....	<b>27</b>
IV. Western Ghats .....	<b>47</b>
V. North East Hills and Brahmaputra Flood Plains .....	<b>63</b>
References .....	<b>75</b>
Acknowledgements & Annexures .....	<b>81</b>



# Executive Summary



©Gurinderjit Singh

Leopards have held cultural significance in ancient civilizations, symbolizing power, agility, and nobility. Today, contemporary portrayals across various mediums continue to highlight their strength, courage, and sensuality. The Indian leopard (*Panthera pardus fusca*) is distributed across a variety of forested habitats in India, Nepal, Bhutan, and parts of Pakistan, excluding mangrove forests and desert habitats (Prater 1980, Daniel 1996). Despite being listed in Schedule I of the Wildlife (Protection) Act, 1972 and classified as “Vulnerable” by the IUCN Red List, leopard populations face significant threats including habitat loss, fragmentation, human-wildlife conflict, poaching, and illegal trade. Leopards occupy a prominent position in the trophic pyramid alongside tigers, lions and dholes exhibiting adaptability in habitat and dietary preferences, and playing a vital role as top predators in a wide array of landscapes across India.

However, their adaptability often leads to conflicts with humans, posing a significant conservation challenge within their range.


Global distribution and population trends indicate significant declines due to various anthropogenic pressures, including habitat loss, prey depletion, and poaching. The NTCA, in collaboration with state forest departments and coordinated by Wildlife Institute of India (WII) conducts comprehensive assessments of tiger habitats, which also include data on leopard density and abundance. Genetic diversity varies across landscapes, with central India exhibiting the highest diversity. Unlike tiger populations, leopard populations show weak structuring across landscapes, with shared genetic affinity between different regions.



©Anurag Nashirabadkar

## Figure E.1:

Leopard abundance, extent and population blocks in India, 2022.



## Methodology

The fifth cycle of leopard population estimation (2022) in India focused on forested habitats within 18 tiger states, covering four major tiger conservation landscapes. Non-forested habitats, arid, and high Himalayas above 2000 msl (~30% area) were not sampled for leopard. To estimate leopard abundance, photo-captures were combined with spatial data on prey, habitat, and anthropogenic factors using a likelihood-based spatially explicit capture mark-recapture (SECR) covariate framework. This method involved three phases:

**Phase I:** Systematic sampling across forested areas within each landscape using M-STriPES Android application and desktop software.

**Phase II:** Utilization of remotely sensed and secondary data to model leopard occupancy and abundance based on habitat characteristics and human impacts.

**Phase III:** Application of SECR models to estimate leopard density, facilitated by camera trap data and individual identification through image and pattern processing software CaTRAT and ExtractCompare respectively. The SECR approach accounted for spatial context and movement patterns, allowing robust population parameter estimates. Covariate-based abundance models were developed for each landscape, with optimal models selected based on statistical criteria. Overall, this comprehensive methodology provided leopard populations across India.

# Countrywide Estimate

The fifth cycle of leopard population assessment (2022) covered 20 states in India, conducting a foot survey spanning 6,41,449 km to estimate carnivore signs and prey abundance.

**Table E.1:** Estimated leopard numbers in each landscape from 2018 and 2022 (Number in parenthesis are one standard error limit of the mean).

State	2018 Population (SE)	2022 Population (SE)
Bihar	98 (8)	86 (3)
Uttarakhand	839 (48)	652 (77)
Uttar Pradesh	316 (39)	371 (54)
Shivalik Hills & Gangetic Plains	1,253 (95)	1,109 (134)
Andhra Pradesh	492 (31)	569 (41)
Telangana	334 (16)	297 (20)
Chhattisgarh	852 (39)	722 (45)
Jharkhand	46 (10)	51 (10)
Madhya Pradesh	3,421 (150)	3,907 (215)
Maharashtra	1,690 (99)	1,985 (122)
Odisha	760 (33)	568 (35)
Rajasthan	476 (39)	721 (112)
Central India & Eastern Ghats	8071 (417)	8,820 (600)
Goa	86 (3)	77 (13)
Karnataka	1,783 (71)	1,879 (261)
Kerala	650 (28)	570 (76)
Tamil Nadu	868 (40)	1,070 (132)
Western Ghats	3387 (142)	3,596 (482)
Arunachal Pradesh	11 (3)	42 (10)
Assam	47 (9)	74 (11)
North Bengal	83 (17)	233 (21)
North Eastern Hills, and Brahmaputra Floodplains	141 (26)	349 (42)
<b>India</b>	<b>12,852 (680)</b>	<b>13,874 (1,258)</b>

Additionally, 3,24,003 habitat plots were sampled for vegetation, human impacts, and ungulate dung. Camera traps were strategically placed at 32,803 locations, resulting in a total of 4,70,81,881 photographs, resulting in 85,488 photo-captures of leopard. The survey comprised of a total effort of 6,41,102 man-days, representing the largest wildlife survey effort worldwide. Leopard presence was recorded in approximately 40% of total 10,846 sampled cells. The esti-

mated leopard population for India was 13,874 (SE 1,258) (Table E.1, Fig E.1). This population adequately covered forested habitat of 13 states barring entire landscape of North East Hills and Brahmaputra Flood Plains where sampling is largely done in Tiger Reserves and partial sampling is done in Uttarakhand (below 2000 msl) and Rajasthan. This estimate of leopard represents ~70% of leopard occupied area. Central India has highest population with Madhya Pradesh having 3,907 individuals followed by Maharashtra, and Karnataka and Tamil Nadu in Western Ghats landscape (Table E.1).

## Shivalik Hills and Gangetic Plains Landscape

Leopard occupy 76% of 351 cells (100 km<sup>2</sup>) which are below 2,000 msl and while numbers have increased in Uttar Pradesh, Uttarakhand has shown a decline in leopard numbers. The leopard population in this landscape is 1,109 (SE 134) (Table E.1, Fig. E.1). There has been a 3.4% decline in overall leopard population in the landscape. Dehradun-Kalsi areas of Uttarakhand were camera trapped for the first time and harbours a sizeable population of leopards (107 minimum leopard number). Leopard number has declined in Ramnagar Forest Division, where tiger numbers have shown a very steep growth in the past four years (Qureshi *et al.*, 2023). Rajaji, Dudhwa and Corbett Tiger Reserves have the largest site wise leopard population (Table E.2) in this landscape. Shivalik landscape has seen an increase in large carnivores and mega-herbivore related conflict in recent years. 65% of the leopard population is present outside Protected Areas in the landscape, which will lead to increase in such conflicts. In the state of Uttarakhand 30% of all the human death and injury cases by wildlife were caused by leopard (Uttarakhand Forest Department, 2024) in the past 5 years. With an increase in tiger number in the landscape (Qureshi *et al.*, 2023), the large carnivore-conflict is what the states need to actively manage. In Uttar Pradesh both leopard and tiger numbers have increased which requires an active engagement for resolution of human-animal conflict by the forest department and civil administration.



©NTCA Tiger Cell

**Table E.2:***The leopard population in Tiger Reserves of India in 2022.*

State	Tiger Reserve	Leopards within Tiger Reserve (SE)	Leopards utilising Tiger Reserve (SE)
<b>Shivalik Hills and Gangetic Plains</b>			
<b>Uttarakhand</b>	Corbett	115 (2)	150 (7)
	Rajaji	171 (6)	215 (10)
<b>Uttar Pradesh</b>	Pilibhit	29 (1)	36 (3)
	Dudhwa	125 (5)	158 (10)
	Ranipur	67(3)	86 (7)
<b>Bihar</b>	Valmiki	78 (1)	86 (3)
<b>Central India and Eastern Ghats</b>			
<b>Andhra Pradesh</b>	Nagarjunasagar Srisailam	270 (5)	360 (14)
<b>Chhattisgarh</b>	Achanakmar	76 (3)	108 (9)
	Indravati <sup>#</sup>	3	3
	Udanti Sitanadi	28 (2)	52 (9)
<b>Jharkhand</b>	Palamau	35 (6)	51 (10)
<b>Madhya Pradesh</b>	Bandhavgarh	146 (4)	176 (8)
	Kanha	157 (2)	209 (9)
	Panna	256 (5)	317 (11)
	Pench	132 (3)	175 (9)
	Sanjay Dubri	110 (3)	154 (9)
	Satpura	215 (2)	256 (8)
<b>Maharashtra</b>	Bor	37 (2)	45 (4)
	Melghat	181 (4)	233 (10)
	Navegaon Nagzira	116 (3)	140 (7)
	Pench	70 (3)	102 (9)
	Sahyadri	87 (2)	135 (10)
	Tadoba Andhari	129 (1)	148 (7)
<b>Odisha</b>	Satkosia	80 (2)	111 (7)
	Similipal	91 (2)	131 (10)
<b>Rajasthan</b>	Mukundara	49 (4)	99 (15)
	Ramgarh Visdhari	19 (1)	25 (3)
	Ranthambore	87 (6)	167 (18)
	Sariska	167 (7)	269 (18)
<b>Telangana</b>	Amrabad	121(2)	173 (10)
	Kawal	19 (1)	25 (3)

State	Tiger Reserve	Leopards within Tiger Reserve (SE)	Leopards utilising Tiger Reserve (SE)
<b>Western Ghats</b>			
<b>Karnataka</b>	Bandipur	138 (3)	185 (9)
	Bhadra	116 (2)	178 (11)
	Biligiri Rangaswamy Temple	66 (3)	98 (9)
	Kali	124 (6)	246 (21)
	Nagarhole	105 (3)	140 (8)
<b>Kerala</b>	Parambikulam	137 (3)	209 (12)
	Periyar	89 (9)	147 (19)
<b>Tamil Nadu</b>	Anamalai**	121 (5)	NA
	Kalakad-Mundanthurai	54 (3)	87 (10)
	Mudumalai	135 (4)	220 (14)
	Sathyamangalam	131 (8)	190 (15)
	Srivilliputhur-Meghamalai	97 (7)	147 (17)
<b>North East Hills and Brahmaputra Flood Plains</b>			
<b>Arunachal Pradesh</b>	Pakke	36 (1)	44 (3)
	Kamlang*	NA	NA
	Namdapha	5	5
<b>Assam</b>	Orang	1	1
	Manas	37 (2)	44 (4)
	Kaziranga	14 (1)	15 (1)
	Nameri	12 (1)	15 (2)
<b>Mizoram</b>	Dampa*	NA	NA
<b>West Bengal</b>	Buxa	61 (1)	74 (4)

\* No evidence of Leopard found during sampling.  
# Minimum identified Leopard from photographs.  
\*\* Estimate derived from covariate model.

## Central India and Eastern Ghats Landscape

The Central India and Eastern Ghats landscape has the largest number of leopards and diverse habitats, from semi-arid regions to mixed deciduous forest. Leopard presence has been identified across all Protected Areas and major forest corridors within the central Indian landscape. While the leopard population in this landscape is growing, largely due to protective measures under the umbrella of tiger conservation, declines are also occurring in states like Chattisgarh, Telangana and Odisha. The surveyed forest in this area has

an estimated total leopard population of 8,820 (SE 600) (Table E.1, Fig E.1). Approximately 68% of the leopard population exists outside the Protected Area. The leopard population estimates for Rajasthan was restricted to current and proposed Tiger Reserves only. Madhya Pradesh holds the largest leopard population in India followed by Maharashtra which signifies that tiger conservation measures also helping the co-predator's recovery (Table E.1). The site wise leopard population is higher in Nagarjunasagar Srisailem, Panna, Sariska, Satpura, Melghat and Kanha Tiger Reserves (Table E.2). The leopard densities are higher in the Tiger Reserves compared to outside Protected Areas, despite the fact that tigers exert regulatory pressure on leopards. Efforts regarding prey recovery and protection need to be strategized and at the same time conflict resolution mechanism need to be evaluated to timely mitigate issues of human-leopard interface.

## Western Ghats Landscape

Although the leopard population in the Western Ghats is widely distributed, it faces the repercussions of habitat loss and fragmentation and poaching (Athreya *et al.*, 2011). The leopard population estimated in the sampled forests of Western Ghats was 3,596 (SE 482) (Tale E.1, Fig. E.1). Leopards in the Western Ghats often inhabited the human-dominated matrices, leopard-human conflict is prevalent in the entire landscape and increased in recent times. While the Nilgiri forests harbour high-density leopard populations (13 leopards/100 km<sup>2</sup>), leopards occur in much lower densities in the scrubland-open forest mosaics of central Karnataka or the evergreen patches of southern Western Ghats (<1 leopard/100 km<sup>2</sup>). In the central and northern Western Ghats, leopard populations are distributed in higher densities inside the Tiger Reserves (Bhadra, Kali, Mudumalai and Sathyamanglam) (Table E.2) while moderate to low densities outside the Protected Areas. The cluster of Anamalai-Parambikulam, located south of the Palghat gap, hosts a population of leopards with good density. The Parambikulam Tiger Reserve, within this cluster, is a hotspot for leopards, where they occur at high densities (Table E.2). In the landscapes of Periyar- Srivilliputhur- Meghamalai and KMTR- Kanyakumari, leopards are found at lower densities, particularly in the evergreen and semi-evergreen forests. More than 65% of the leopard population is present outside the Protected Areas in the Western Ghats landscape. Although the leopard population in the landscape showed a stable trend in the consistent sampling area, in certain areas, the leopard population declined significantly, e.g., the Wayanad landscape, which needs to be appropriately examined. At present, leopards often use the mosaics of forests, plantations, and human land uses outside the Protected Areas. However, the situation is changing as people start fencing their lands in order to reduce crop damage, and mushrooming developmental projects have come up, which are causing fragmentation in the population.


## North East Hills and Brahmaputra Floodplains Landscape

North East Hills and Brahmaputra Flood Plains is a mosaic of forested habitat interspersed with human habitation and land cover patterns which provides ample cover for adaptable species like leopard. Although leopard presence is evident across the landscape, population estimates of leopards are available only from the Tiger Reserves, and some areas outside Reserves in North Bengal. Despite efforts put in since the last five cycles of All India Tiger Estimation, large forested areas of Assam and Arunachal Pradesh outside Tiger Reserves were not sampled. The sampling remains inadequate in this landscape and needs attention to expand the scope of adequate monitoring. A total of 15 sites were sampled in this landscape in camera trap based mark-recapture framework and leopards were photo-captured in 11 sites. Estimated leopard population of this landscape is 349 (SE 42). Although there is an increase in the leopard population as compared to earlier cycle, this is mostly because of the increase in sampling effort in North Bengal *Dooars*. Growing human leopard conflict in this landscape owing to habitat mosaic and developmental projects is a major concern towards leopard conservation.

## Conservation Implications

The population growth in common area from 2018 to 2022 across India was 1.08 %/annum, Shivalik Hills and Gangetic Plains recorded -3.4%/annum decline, while Central India and Eastern Ghats, Western Ghats and North East Hills and Brahmaputra Flood Plains recorded growth of 1.5%, 1.0% and 1.3% per annum respectively. The population in last four years is stable, which also indicates the growth is minimal and in comparison to tiger its population is likely getting impacted by people in multiple use area. Current trend of poaching is unknown but seems to be likely cause of stable population, which includes commercial poaching and people's retaliation to conflict with leopards.







# I. Status of leopards in India

Qamar Qureshi, Yadvendradev V. Jhala, Vishnupriya Kolipakam, Satya Prakash Yadav, Ujjwal Kumar, Shikha Bisht, Ayan Sadhu, Deb Ranjan Laha, Swati Saini, Rajendra Garawad and Kausik Banerjee

Leopards commanded great reverence in ancient cultures (Egyptian, African and Indian cultures), symbolizing traits of power, agility, and nobility. Among Native cultures, the leopard's hunting prowess and camouflage were deeply respected, with tales highlighting their significance in transformational narratives and superstitions. Myths and folklore surrounding leopards often depict them as bold, cunning spirits intertwined with the spirit realm. Today, contemporary portrayals of leopards across fashion, literature, cinema, and art persist in symbolizing qualities of strength, courage, and sensuality (<https://roaringrealms.com/leopard-myths-cultural-significance/>). The Indian leopard, with its spotted coat and powerful presence, has long held a significant place in Indian art and culture. More than just a magnificent predator, it has been woven into the fabric of myths, folklore, and artistic expressions, representing a captivating blend of fear, power, and beauty.

The Indian leopard (*Panthera pardus fusca*) is found in a variety of forested habitats, including tropical rainforests, dry deciduous forests, temperate forests, and coniferous forests. The subspecies is found in India, Nepal, Bhutan, and parts of Pakistan, and it is not found in mangrove forests or desert habitats (Prater 1980, Daniel 1996). According to the Wildlife (Protection) Act, 1972 leopard is listed in Schedule I, which accorded highest protection and as per International Union for Conservation of Nature (IUCN) Red List of Threatened Species, it is classified as "Vulnerable". Its population is decreasing across its range, primarily due to habitat loss, fragmentation, human-wildlife conflict, poaching, and illegal trade (Athreya *et al.*, 2011 and Raza *et al.*, 2012).

Following tigers (*Panthera tigris*), dholes (*Cuon alpinus*) and lions (*Panthera leo*), leopards hold a prominent position in the trophic pyramid. While tigers migrated to India from the East, specifically the Malayan realm, leopards share an origin similar with lions, originating in the Ethiopian realm. Interestingly, leopards entered India through the Western corridor, preceding the arrival of lions and tigers (Mani 1974). In the Himalayas, leopards are occasionally sympatric with snow leopards (*Panthera uncia*) and have been sighted at altitudes as high as 5,200 meters (Uphyrkina *et al.*, 2001). Leopards exhibit remark-

able adaptability in terms of habitat and dietary preferences, thriving in agro-pastoral areas, plantations, and even near human settlements, both rural and urban (Nowell and Jackson, 1996). Known for their prolific breeding rates and faster life history traits compared to tigers, leopards have demonstrated an annual growth rate of 15% in parts of Kanha Tiger Reserve in Central India (Kumar *et al.*, 2019).

Being one of the most widely distributed felids, leopards owe their extensive range to their dietary flexibility and ability to inhabit various environments, including human-dominated landscapes (Sunquist and Sunquist, 2002; Hayward *et al.*, 2006; Athreya *et al.*, 2013; Gubbi *et al.*, 2020). However, this adaptability often leads to conflicts with humans, presenting a significant conservation challenge within their range (Rahalkar, 2008; Athreya *et al.*, 2011; Navya *et al.*, 2014; Sidhu *et al.*, 2017; Naha *et al.*, 2018).

The global distribution and population of leopards have witnessed significant declines due to habitat loss, depletion of prey species, human-wildlife conflicts, and poaching over the past century. Recent studies suggest a range loss of 48–67% for leopards in Africa and 83–87% in Asia (Jacobson *et al.*, 2016), corroborating genetic evidence indicating a 75–90% human-induced population decline within the last ~120–200 years in India (Bhatt *et al.*, 2020).

The National Tiger Conservation Authority (NTCA), in collaboration with state forest departments, conservation NGOs, and coordinated by the Wildlife Institute of India, conducts a comprehensive national assessment of "Tigers, Co-predators, Prey, and their Habitat" every four years since 2006. While past assessments primarily focused on occupancy for species other than tigers and leopards, the 2014 cycle marked the first reporting of leopard density and abundance across tiger habitats in India, estimating the leopard population at 7,910 (SE 1,344) and 12,852 (SE 680) individuals in 2014 and 2018 (Jhala *et al.*, 2015 & Jhala *et al.*, 2021) respectively. This report presents the status of leopards based on camera trap data and occupancy surveys conducted in 2022 across 20 states of India where tigers are present, utilizing a spatially explicit capture-recapture framework. Given the substantial coverage of forested landscapes in these 18 states using camera traps, photo-capture

data are also employed to reliably estimate occupancy and abundance. The genetic diversity of leopard populations in India varies across landscapes, with Central India exhibiting the highest diversity. Unlike tiger populations, leopard populations show weak structuring across landscapes, as indicated by genetic analysis (Jhala *et al.*, 2019). The data reveals three clusters, with largely overlapping signatures. While there are no genetically distinct populations observed at lower levels of division, Terai and parts of Central Indian leopards appear distinct from Western Ghats and North Eastern populations. There is a shared affinity of genes between Eastern Ghats, Western Ghats and North Bengal populations. Notably, leopard populations exhibit genetic continuity throughout the Western Ghats, contrasting with the structured nature of tiger populations in the same regions (Kolipakam *et al.*, 2019).

Conservation efforts for the Indian leopard require habitat restoration initiatives, protection outside Protected Area system and the implementation of measures to mitigate human-leopard conflict. There is a need to collaborate with local communities to raise awareness, and promote coexistence between humans and leopards. Research on leopard ecology, behavior, and population dynamics is needed for informed conservation strategies and management decisions.

## Methodology

The leopard population estimation was carried out within forested habitats in tiger occupied states of the country. Four major tiger conservation landscapes, 1) Shivalik Hills and Gangetic Plains, 2) Central India and Eastern Ghats, 3) Western Ghats and, 4) North East Hills and Brahmaputra Flood Plains, were sampled for estimating leopard abundance. Details of these landscapes are available in Qureshi *et al.*, 2023. Non-forested habitats like commercial plantations, and other land use forms, higher elevations in Himalayas (>2000 msl), arid landscapes and majority of North East landscapes, where leopards are known to occur, were not sampled. Therefore, the current estimate of leopards presented here should be considered as ~70% of leopard's landscape sampled.

To arrive at the abundance of leopards, photo-captures of individual leopards were used in combination with spatial data on prey, habitat, and anthropogenic factors as covariates in a likelihood based spatially explicit capture mark-recapture (SECR) covariate framework (Efford 2015). This method entails estimating spatial covariates of relative abundance of leopard and other co-predators, ungulates, human impact indices, and habitat characteristics across all potential tiger habitats in India, at a fine spatial resolution of a forest beat which is on average about 15 km<sup>2</sup> (Phase I and II). Within each landscape, sampling was carried out at a high density, and one double sided camera station placed in a 2 km<sup>2</sup> area (Phase III) (Fig. I.1). The raw counts of abundance obtained from the sample space are calibrated

against absolute density obtained from the areas where camera trapping is carried out. In this SECR approach, spatial information on capture-mark-recapture (that accounts for detection correction) is used with spatial covariates of leopard sign intensity, prey abundance, human disturbance and habitat characteristics to estimate leopard population directly in the camera trapped area, and extrapolates to forested areas with leopards but not camera trapped, based on covariates.

### Phase I

Sampling is systematically carried out across all forested areas within each landscape. The average size of a forest beat in India is 15-16 Km<sup>2</sup>. These spatial administrative units are the basis of ensuring fine scale sampling across the forested habitats in each tiger bearing landscape. To ensure appropriate data collection, the State Forest Departments were trained by NTCA-WII Tiger Cell in the Phase I sampling protocols to be carried out across current and potential tiger habitats (in Tiger Reserves, Protected Areas, Reserve Forests and in all Wildlife and Territorial divisions) across 20 states (Figure I.2). Data was recorded digitally using M-STriPES (Monitoring System for Tigers: Intensive Protection and Ecological Status) ecological android mobile application. The protocol for Phase I and the details of the forms are explained in Qureshi *et al.*, 2023. Broadly, the data is collected on carnivore sign encounters (Form 1), Ungulate abundance (Form 2), Habitat status (Form 3A, 3B, 3C) and index of herbivore presence (Form 4) (Fig I.1) (Table I.1).

Since data is collected in a digital format on the M-STriPES mobile Android app, it was directly imported and analysed on the complementing desktop software. Phase I data was received from 628 Forest Divisions of India and these were processed using M-STriPES desktop software. Data for each spatial and temporal replicate was recorded at the beat scale (occupancy surveys, line transects, and plots) were transferred to the standard 25 and 100 km<sup>2</sup> cells for analysis and subsequent inference. Tiger and leopard sign encounter rates, ungulate encounter (direct sighting) rates, ungulate dung density, human disturbance indices (signs of livestock, human trails, wood cutting, lopping, grass removal) were computed as average encounter rates for 5x5 and 10x10 km cells based on effort (km of survey) invested in each cell.

### Phase II

Remotely sensed data is used to obtain covariates like habitat characteristics and anthropogenic impacts that are likely to determine the distribution and abundance of leopards. This data is used then to model leopard occupancy and abundance. Habitat characteristics were surrogated by forest area, vegetation cover [Normalized Difference Vegetation Index, (NDVI)], forest patch size, forest core areas, elevation, distance from Protected Areas and drainage density. Human impacts were surrogated by human footprint, distance to night lights, night light intensity, distance to roads and density of road network (Fig. I.1, Table I.2).

**Table I.1:**

*Country wide sampling effort for ground surveys and camera traps during Phase I and III of each state and landscape, 2022.*

State	Number of Trails	Total Length of Trails (Kms)	Number of Transects	Total Length of Transects (Kms)	Number of Habitat Plots	Camera Trap Numbers
Bihar	232	1775	499	981	1372	429
Uttar Pradesh	1270	6412	1237	2408	4396	1343
Uttarakhand	2735	12389	2522	4299	8256	2165
Shivalik Hills and Gangetic Plains	4237	20576	4258	7688	14024	3937
Andhra Pradesh	3456	16597	3430	6637	11090	989
Chhattisgarh	9855	46176	8422	16185	31664	459
Jharkhand	976	4600	732	1457	4570	323
Madhya Pradesh	26757	139651	26341	54256	96924	6894
Maharashtra	16331	78016	16124	31210	56512	4872
Odisha	9623	52633	9531	19522	33544	733
Rajasthan	988	4449	981	1911	3422	685
Telangana	6633	29188	5599	10502	18264	1578
Central India and Eastern Ghats	74619	371310	71160	141680	255990	16533
Karnataka	8874	45323	10002	18297	31742	5163
Kerala	1522	7463	1201	2361	4156	1314
Tamil Nadu	1892	11019	1845	3681	5376	3650
Goa	174	765	165	324	526	95
Western Ghats	12462	64570	13213	24663	41800	10222
Assam	389	1816	483	856	1356	619
North Bengal	786	4011	586	1203	2174	329
Mizoram	87	330	NA	NA	8064	43
Arunachal Pradesh	309	840	NA	NA		407
Nagaland	178	568	NA	NA		NA
North East Hills and Brahmaputra Flood Plains	1749	7565	1069	2059	11594	1398
Sundarbans	315	1339	NA	NA	595	713
<b>India</b>	<b>93,382</b>	<b>4,65,360</b>	<b>89,700</b>	<b>1,76,090</b>	<b>3,24,003</b>	<b>32,803</b>

**Table I.2:**

Remotely sensed spatial data based on field sampling and secondary data used for modelling occupancy and abundance of leopards.

S. No	Dataset	Source	Spatial Resolution
1	Leopard sign encounter rate	Phase I survey, AITE 2022	25 km <sup>2</sup>
2	Tiger sign encounter rate	Phase I survey, AITE 2022	25 km <sup>2</sup>
3	Prey encounter rate (Chital, Sambar, Wild Pig, Barking Deer, Langur)	Phase I survey, AITE 2022	25 km <sup>2*</sup>
4	Prey faecal pellet (Chital, Sambar, Wild Pig, Barking Deer, Langur)	Phase I survey, AITE 2022	25 km <sup>2*</sup>
5	Extractive human use (Wood cutting, lopping, human & livestock trails, people seen)	Phase I survey, AITE 2022	25 km <sup>2*</sup>
<b>Remotely sensed data</b>			
3	Night time lights Intensity (2021)	C. D. Elvidge, K. Baugh, M. Zhizhin, F. C. Hsu, and T. Ghosh, "VIIRS night-time lights," <i>International Journal of Remote Sensing</i> , vol. 38, pp. 5860–5879, 2017.	500 m
4	Normalised Difference Vegetation Index-Pre and post monsoon	Vermote, E., Justice, C., Claverie, M., & Franch, B. (2016). Preliminary analysis of the performance of the Landsat 8/OLI land surface reflectance product. <i>Remote Sensing of Environment</i> , 185, 46-56.	30 m
5	Digital Elevation Model	NASA Shuttle Radar Topography Mission (SRTM) (2013). Shuttle Radar Topography Mission (SRTM) Global. Distributed by OpenTopography. <a href="https://doi.org/10.5069/G9445JDF">https://doi.org/10.5069/G9445JDF</a> .	30m
6	Global human modification Index (2016)	Kennedy, C. M., J. R. Oakleaf, D. M. Theobald, S. Baruch-Mordo, and J. Kiesecker. 2020. Global Human Modification of Terrestrial Systems. Palisades, NY: NASA Socioeconomic Data and Applications Center (SEDAC). <a href="https://doi.org/10.7927/edbc-3z60">https://doi.org/10.7927/edbc-3z60</a> .	1000 m
7	Forest Cover (2017)	India State of Forest Report (2017). Forest Survey of India, Ministry of Environment, Forest and Climate Change, Government of India	23.5 m
8	Ruggedness	Derived from Digital Elevation Model	30 m
9	Distance to Nightlights	Euclidean distance from Nightlights data	1000 m
10	Distance to water sources	Euclidean distance derived from global surface water dataset (Pekel <i>et al.</i> , 2016)	1000 m
11	Distance to Protected Areas	Data archived from Wildlife Database Cell, WII and Project Tiger database	1000 m

\*Data collected on 2-3 km line transects & plots laid at every 400 m were extracted at 25 km<sup>2</sup> cell

### Phase III

Spatially Explicit Capture Recapture (SECR) models integrate the spatial context of capturing and recapturing individuals with their temporal capture history to estimate density. By linking the detection process to the actual space utilization of an animal, SECR provides robust population

parameter estimates (Borchers and Efford, 2008). Individual identification and abundance estimation of tigers and leopards are facilitated by their distinctive stripes and rosettes, within the framework of capture-mark-recapture. Camera traps were placed at 32,803 locations, spread across 175 sites for mark recapture analysis (Figure I.2, I.3).

Camera traps were systematically positioned in the sampling area through a 2 km<sup>2</sup> cell overlay, deploying at least one pair of cameras in each cell. The cameras were strategically located to maximize photo-captures of tigers and leopards, identified through thorough sign surveys. Each cell received a unique code and was positioned within the nationwide 100 km<sup>2</sup> cell established since the initial National Tiger Status Estimation cycle in 2006, ensuring consistent spatial comparison for subsequent inferences. Sampling occurred concurrently in a minimum block of 200 km<sup>2</sup>, expanding to larger blocks if more camera traps were available to cover areas exceeding 200 km<sup>2</sup>. A minimum spacing of approximately 1 km between camera trap locations was maintained. Typically, cameras operated for 25 to 35 days at each site, with an average effort exceeding 1,200 trap-nights per ~100 km<sup>2</sup>.

**Phase III analysis:** As explained in Qureshi *et al.*, 2023, in collaboration with the Indraprastha Institute of Information Technology, New Delhi, an AI-based image processing tool was developed to automatically geotag and categorize camera trap images by species (Fig. I.1, I.3 & I.4). This image processing software, known as CaTRAT (Camera Trap Data Repository and Analysis Tool) (Cheema *et al.*, 2018), was employed for geotagging, coding, and organizing images into individual species folders. The geo-tagged images underwent careful examination to identify potential software misclassifications, and the separated photos of tigers and leopards were further processed for individual identification of these species.


Leopard images were grouped utilizing Hotspotter (Crall *et al.*, 2013), and final leopard identification was conducted using ExtractCompare (Hiby *et al.*, 2009). A total of 85,488 Leopard photographs were collected from camera traps. In ExtractCompare, a three-dimensional surface model of a leopard is overlaid on a leopard photo to accommodate pitch and roll associated with body posture before extracting the spot pattern (Figure I.5). Employing an automated process, pattern recognition software searches the image database to calculate similarity scores between digitized leopard coat patterns, facilitating the recognition of common and unique individuals. Initially, leopard(s) photographed at each camera trap site were identified using this method. Subsequently, comparisons were made between leopard photographs from adjacent sites and within each landscape using the National database to eliminate any duplicate leopards and comprehend leopard dispersal events. Once individual leopards were identified, a spatial capture history matrix for each leopard was developed for each site, including camera trap IDs, their coordinates, and the deployment and operation history of each camera.

We utilized likelihood-based SECR methods (Borchers *et al.*, 2008; Efford, 2011) to estimate leopard abundance from camera trap data. The two fundamental detection parameters in SECR include the detection probability ( $g_0$ ) at the home range center of the animal and a parameter for spatial movement ( $\sigma$ ). A habitat mask, incorporating a realistic buffer around the camera trap array to exclude non-habitat, was provided in our analysis. Density was modeled as a function of covariates, including tiger and leopard sign encounter rates, prey encounter or dung densities, and human footprint variables from ground surveys and remotely sensed data. These covariates were employed within SECR to model leopard density through the secr package (Efford, 2015) in the R programming environment (R Core Team, 2023). Covariate-based abundance models were developed for each landscape to estimate leopard abundance within tiger-occupied forests. The optimal covariate model was selected based on Akaike Information Criteria (Akaike, 2011) for each landscape. In areas where leopards were detected but not camera-trapped, their numbers were estimated by predicting leopard density from covariates (prey, habitat, and human disturbances) using the best model or model-averaged parameters.

We used a combination of several covariates (including tiger and leopard sign encounters, extractive use indices, NDVI, elevation, human-modified indices, and distance to Protected Areas) in a spatially explicit capture-recapture model to predict leopard density outside the camera-trapped area (except North East Hills and Brahmaputra Flood Plains where sufficient Phase I information was not available for areas outside camera trapped sites). The selection of the most parsimonious covariate model for each landscape was based on the lowest AICc value.

For the Shivalik Hills and Gangetic Plains landscapes, the best model included leopard sign encounters ( $\beta = 0.11 \pm 0.03$ , positively correlated) and tiger sign encounters ( $\beta = -0.25 \pm 0.06$ , negatively correlated). In the Central India and Eastern Ghats landscape, the optimal model comprised leopard sign encounters ( $\beta = 0.16 \pm 0.01$ , positively correlated) and prey encounters (chital, sambar, barking deer, langur, and wild pig) ( $\beta = 0.26 \pm 0.03$ , positively correlated). For the Western Ghats landscape, the preferred model incorporated leopard sign encounters ( $\beta = 0.25 \pm 0.02$ , positively correlated), prey encounters (chital, sambar, barking deer, langur, and wild pig) ( $\beta = 0.09 \pm 0.01$ , positively correlated), and extractive use from the forest ( $\beta = -0.07 \pm 0.02$ , negatively correlated). The selection of these models was based on the lowest AICc values.


**Figure I.1:**  
Outline of Sampling framework for leopard estimation






**Figure 1.2.**

*Spatial coverage of sampled forests for carnivore signs where leopard evidence was recorded is shown in orange and black denotes camera trap locations with leopard photo-captures in India, 2022.*




**Figure 1.3.**

*Workflow of species identification from camera trap images using artificial intelligence based tool, CaTRAT*



## Figure 1.4.

Process of individual identification of leopards using Program ExtractCompare



# Abundance Assessment of Leopards in India

The fifth cycle of leopard population (2022) was conducted in 20 states of India, encompassing a foot survey of 6,41,449 km for carnivore signs and prey abundance estimation. Additionally, 3,24,003 habitat plots were sampled for vegetation, human impacts, and ungulate dung. Camera traps were deployed at 32,803 locations, resulting in 4,70,81,881 photographs, of which 85,488 were of leopards. The total effort invested in the survey was 6,41,102 man-days, making it the world's largest effort invested in any wildlife survey to date. Leopard presence was recorded in 40% (4321) of sampled cells (n=10846). The leopard estimate for India was 13,874 (SE 1,258) and 67% of this leopard population is outside Protected Areas. This estimate of leopard represents ~70% of leopard occupied area in 18 states.

**Table I.3:** Estimated leopard numbers in each landscape from 2018 and 2022 (Number in parenthesis are one standard error limits of the mean).

State	2018 Population (SE)	2022 Population (SE)
Bihar	98 (8)	86 (3)
Uttarakhand	839 (48)	652 (77)
Uttar Pradesh	316 (39)	371 (54)
Shivalik Hills and Gangetic Plains	1,253 (95)	1,109 (134)
Andhra Pradesh	492 (31)	569 (41)
Telangana	334 (16)	297 (20)
Chhattisgarh	852 (39)	722 (45)
Jharkhand	46 (10)	51 (10)
Madhya Pradesh	3,421 (150)	3,907 (215)
Maharashtra	1,690 (99)	1,985 (122)
Odisha	760 (33)	568 (35)
Rajasthan	476 (39)	721 (112)
Central India and Eastern Ghats	8071 (417)	8,820 (600)
Goa	86 (3)	77 (13)
Karnataka	1,783 (71)	1,879 (261)
Kerala	650 (28)	570 (76)
Tamil Nadu	868 (40)	1,070 (132)
Western Ghats	3,387 (142)	3,596 (482)
Arunachal Pradesh	11 (3)	42 (10)
Assam	47 (9)	74 (11)
North Bengal	83 (17)	233 (21)
North East Hills and Brahmaputra Flood Plains	141 (26)	349 (42)
India	12,852 (680)	13,874 (1,258)

This population adequately covered forested habitat of 13 states barring states of North-Eastern Hills and Brahmaputra Flood Plains where sampling is largely done in Tiger Reserves and partial sampling is done in Uttarakhand (below 2000 msl) and Rajasthan (Protected Areas).

Central India has highest population with 8,820 (SE 600) leopards followed by Western Ghats 3,596 (SE 482) and Shivalik Hills and Gangetic Plains having 1,109 (SE 134) individuals. The partial sampling of North East Hills and Brahmaputra Flood Plains have population of 349 (42) leopards. The states which have largest population of leopards were Madhya Pradesh (3,907), Maharashtra (1,985), Karnataka (1,879) and Tamil Nadu (1,070) (Table I.3) and Tiger Reserves or sites with highest leopard population are, Nagarajunasagar Srisailam, Panna, Satpura, Sariska, Melghat, Kali, Mudumalai, Kanha and Parambikulam (Table I.4).

The population growth in common area from 2018 to 2022 across India was 1.08%/annum, Shivalik Hills and Gangetic Plains recorded -3.4%/annum decline, while Central India and Eastern Ghats, Western Ghats and North East Hills and Brahmaputra Flood Plains recorded growth of 1.5%, 1.0% and 1.3% per annum respectively.


In the Shivalik Hills Landscape, it is crucial to monitor tiger and leopard dynamics annually, not only within Tiger Reserves but also in territorial and forest divisions, especially in Uttarakhand where a significant portion of the leopard population (79%) resides outside Protected Areas while for entire landscape it is 65%. The Rajaji, Dudhwa and Corbett Tiger Reserves have substantial leopard population (Table I.4). Annual camera trapping similar to Phase IV monitoring is recommended to track leopard populations effectively. Additionally, consideration should be given to declaring areas like Terai East and Nandhaur Wildlife Sanctuary as Tiger Reserves to facilitate staff training and annual monitoring efforts.

Nagarajunasagar Srisailam, Panna, Sariska and Satpura Tiger Reserves have very good leopard population (Table I.4). In Central India, a substantial portion (68%) of the leopard population inhabits human-use forests outside Protected Areas, increasing vulnerability to poaching and human-leopard conflicts. This region is a hotspot for leopard poaching (Raza *et al.*, 2012), necessitating strategic planning to address challenges such as prey recovery, mitigating road-hit mortalities, and capacity building for managing human-leopard conflicts. Effective patrolling and law enforcement are particularly critical in states like Jharkhand, Chhattisgarh, Odisha, and Telangana.

In the Western Ghats, leopard populations exhibit stability, but 65% of leopards reside outside Protected Areas, raising concerns for conservation efforts. Substantial population of leopards occur in Kali, Parambikulam and Mudumalai (Table I.4). This scenario increases the likelihood of leopard-human conflicts, leading to retaliatory killings of

## Figure 1.5

Leopard density, extent and population blocks in India, 2022.



leopards, especially in states like Karnataka, Kerala, Tamil Nadu, and Goa. Landscape-level plans are imperative to study conflict dynamics, and long-term studies focusing on leopard population dynamics and behavior are essential for comprehensive understanding. These measures can contribute to the development of a landscape-level leopard conservation plan for the Western Ghats.

North East Hills and Brahmaputra Flood Plains comprise a mosaic of forested habitats interspersed with human habitation and various land cover patterns, providing ample cover for adaptable species like leopards. While leopard presence is evident across the landscape, population estimates are only available from Tiger Reserves and some areas outside Reserves in North Bengal. Despite efforts made in the past

five cycles of the All India Tiger Estimation, large forested areas of Assam and Arunachal Pradesh outside tiger reserves have remained unsampled, indicating inadequate sampling in this landscape and necessitating attention to expand monitoring scope. A total of 15 sites were sampled in this landscape using a camera trap-based mark-recapture framework, with leopards being photo-captured in 11 sites. The estimated leopard population in this landscape is 349 (SE 42). Although there has been an increase in the leopard population compared to earlier north Bengal *Doors*. The growing human-leopard conflict in this landscape, stemming from habitat mosaic and developmental projects, poses a significant concern for leopard conservation

**Table I.4:***Leopard population in Tiger Reserves of India in 2022.*

State	Tiger Reserve	Leopards with- in Tiger Reserve (SE)	Leopards utilising Tiger Reserve (SE)
<b>Shivalik Hills and Gangetic Plains</b>			
<b>Uttrakhand</b>	Corbett	115 (2)	150 (7)
	Rajaji	171 (6)	215 (10)
<b>Uttar Pradesh</b>	Pilibhit	29 (1)	36 (3)
	Dudhwa	125 (5)	158 (10)
	Ranipur	67(3)	86 (7)
<b>Bihar</b>	Valmiki	78 (1)	86 (3)
<b>Central India and Eastern Ghats</b>			
<b>Andhra Pradesh</b>	Nagarjunasagar Srisaïlam	270 (5)	360 (14)
<b>Chhattisgarh</b>	Achanakmar	76 (3)	108 (9)
	Indravati <sup>#</sup>	3	3
	Udanti Sitanadi	28 (2)	52 (9)
<b>Jharkhand</b>	Palamau	35 (6)	51 (10)
<b>Madhya Pradesh</b>	Bandhavgarh	146 (4)	176 (8)
	Kanha	157 (2)	209 (9)
	Panna	256 (5)	317 (11)
	Pench	132 (3)	175 (9)
	Sanjay Dubri	110 (3)	154 (9)
	Satpura	215 (2)	256 (8)
<b>Maharashtra</b>	Bor	37 (2)	45 (4)
	Melghat	181 (4)	233 (10)
	Navegaon Nagzira	116 (3)	140 (7)
	Pench	70 (3)	102 (9)
	Sahyadri	87 (2)	135 (10)
	Tadoba Andhari	129 (1)	148 (7)
<b>Odisha</b>	Satkosia	80 (2)	111 (7)
	Similipal	91 (2)	131 (10)
<b>Rajasthan</b>	Mukundara	49 (4)	99 (15)
	Ramgarh Visdhari	19 (1)	25 (3)
	Ranthambore	87 (6)	167 (18)
	Sariska	167 (7)	269 (18)
<b>Telangana</b>	Amrabad	121 (2)	173 (10)
	Kawal	19 (1)	25 (3)
<b>Western Ghats</b>			
<b>Karnataka</b>	Bandipur	138 (3)	185 (9)
	Bhadra	116 (2)	178 (11)
	Biligiri Rangaswamy Temple	66 (3)	98 (9)
	Kali	124 (6)	246 (21)
	Nagarhole	105 (3)	140 (8)

State	Tiger Reserve	Leopards with- in Tiger Reserve (SE)	Leopards utilising Tiger Reserve (SE)
Kerala	Parambikulam	137 (3)	209 (12)
	Periyar	89 (9)	147 (19)
Tamil Nadu	Anamalai**	121 (5)	NA
	Kalakad-Mundanthurai	54 (3)	87 (10)
	Mudumalai	135 (4)	220 (14)
	Sathyamangalam	131 (8)	190 (15)
	Srivilliputhur-Meghamalai	97 (7)	147 (17)
<b>North East Hills and Brahmaputra Flood Plains</b>			
Arunachal Pradesh	Pakke	36 (1)	44 (3)
	Kamlang*	NA	NA
	Namdapha	5	5
Assam	Orang	1	1
	Manas	37 (2)	44 (4)
	Kaziranga	14 (1)	15 (1)
	Nameri	12 (1)	15 (2)
Mizoram	Dampa*	NA	NA
West Bengal	Buxa	61 (1)	74 (4)

\* No evidence of Leopard found during sampling.  
# Minimum identified Leopard from photographs.  
\*\* Estimate derived from covariate model.

A study conducted in 2012 suggests an alarming rate of poaching, 4 leopards are being poached every week in the past decades (Raza *et al.*, 2012). These figures may represent only a fraction of the actual poaching and trade in leopard parts occurring in India. Poaching of large carnivores may alter demography, behaviour of the target species also leaves cascading impact on the native biodiversity. Beyond poaching, habitat fragmentation due to developmental activities such as the development of linear infrastructure and mining poses significant threats to leopards. Mineral mining, linear development without appropriate mitigation that intersect its biodiverse forest habitats. Over the last century, leopards in India have experienced a substantial human-induced population decline (Bhatt *et al.*, 2020).


With increase in human populations and encroachments further into leopard habitats, incidents of human-leopard conflict are on the rise. In some instances, these conflicts escalate into violence, with communities retaliating against leopards that have preyed on their livestock or injures human. While recent studies have accumulated substantial knowledge about leopard populations, habitat use, food habits, and human-leopard conflict, there remains a notable gap in understanding their behaviour in varying natural conditions. With the leopard population expected to increase in the future, a concerted effort toward long-term monitoring, coupled with a deeper understanding of their natural history and behaviour, is imperative to devise effective management strategies.



©Anuradha Marwah

## Figure I.6:

Leopard and Tiger abundance and distribution across India, 2022



The three major carnivores in the tropical forests of India are tigers, leopards, and dholes. The dominance hierarchy is Tiger > Dholes > Leopard (Srivathsa *et al.*, 2023). It is important to understand the distribution, abundance, and interaction amongst them, as they play an important role in shaping the community structure in tropical forests. Tiger and dhole have almost similar distribution except in Shiwalik Hills, Gangetic Plains, and Sunderbans. In their former habitat, dholes were persecuted and are almost wiped out of this area, and Sunderbans is not a dhole habitat. The leopard is more widespread across India in almost all habitats except deserts and Sunderbans. The tigers are distributed in forest ecosystems that are less disturbed and have good deer density. The principal prey of all three carnivores are chital, sambar, hog deer, barking deer, wild pigs, and primates (Johnsingh 1992; Karanth 1995; Majumdar 2013; Steinmetz *et al.*, 2023). The three species exhibit spatio-temporal interaction, behavioral mechanisms, and prey diversity that allow for a form of co-occurrence, despite some level of avoidance and occasional aggression. There are complex interactions and dynamics within large carnivore guilds, as

well as the influence of human activities on these interactions (Athreya, 2012; Gubbi *et al.*, 2017; Jhala *et al.*, 2020 & 2021; Srivathsa *et al.*, 2023).

The countrywide data indicates a positive or no relationship between the tiger and leopard populations in most of the areas except parts of the Gangetic and Brahmaputra floodplains (Fig I.6). Terrain complexity provides space for leopards and other large carnivores to persist with spatio-temporal segregation (Kumar *et al.*, 2019). The extractive use by humans has a negative relationship, indicating the interaction of protected areas providing prey and protection, as most of the high-density multi-carnivore areas are tiger reserves or other protected areas. The grassland management in protected areas at village-relocated sites has provided an opportunity for the sustenance of large prey biomass. It is important to maintain the grasslands and habitat free from invasive plant species, and the removal of ungulates on a large scale should be avoided as large-medium carnivore guilds coexist in a productive ecosystem.







# II. Shivalik Hills and Gangetic Plains Landscape

Shikha Bisht, Ujjwal Kumar, Vishnupriya Kolipakam, Swati Saini, Vaishnavi Gusain, Monika Saraswat, Hemant Singh, Satya P. Yadav, Yadvendradev V. Jhala and Qamar Qureshi



©Harsh Sethi

## Shivalik Hills and Gangetic Plains Landscape

The Shivalik Hills and Gangetic Plains landscape in India spans across the states of Himachal Pradesh, Uttarakhand, Uttar Pradesh and Bihar is comprised of three parallel geological zones, viz. the Shivaliks, the Bhabar tract and the Terai plains. We assessed leopard population only in the potential areas where tiger could occur in this landscape. Leopards are distributed in the Shivalik Forest Division adjoining Rajaji Tiger Reserve, towards Himachal Pradesh, and in the higher hilly regions of the state of Uttarakhand. But these areas were outside the purview of the current project's study area and hence, we provide the estimate of leopard population in potential tiger bearing forests below 2000 msl.


Leopards are distributed across the Shivalik Hills and Gangetic Plains, and are reported to use non-forested areas that include vicinity of human habitations, plantations and agricultural fields (Fig. II.1). Major wild prey for leopard in the Rajaji Tiger Reserve was chital, sambar, hog deer, barking


deer and wild pig. Patel *et al.*, 2023 reported that in Shivalik and Terai region, diet of leopards was primarily composed of large and medium-sized prey species (85%), whereas small prey species made up just 15%. Sambar (23%) and Chital (45 %) had the majority of contribution to the diet. In all, nine different species of prey were found in the diet of leopard, large-bodied Sambar, Nilgai, and Livestock; medium-bodied Chital, Wild pig, and Hog deer; and small-bodied Langur, Hare, and Peafowl. Most of the studies in the landscape focus on human-leopard conflict (Naha *et al.*, 2018 and 2020).



©Gurinderjit Singh

**Figure II.1:** Change in leopard distribution in Shivalik Hills and Gangetic Plains from 2018 to 2022.





**Figure 11.2:** Spatially explicit leopard density modelled from camera traps-based capture-mark-recapture and covariates of leopard sign for Shivalik Hills and Gangetic Plains Landscape 2022.

# Leopard Occupancy, Population Extent and Abundance

Phase I data collected by the forest department and camera trap based photo-captures show leopard to be distributed across the forested areas of Uttarakhand, Uttar Pradesh and parts of Bihar. The population in much of Uttar Pradesh is contiguous with habitats in Nepal. Leopards were also reported from the higher reaches of Himalayas (Narendranagar, Almora, Nainital and Champawat forest division) wherever the habitat was sampled.

Of the 351 cells of 100 km<sup>2</sup> that were sampled, 268 cells (76 %) were occupied by leopards in the landscape. The landscape sampling has saturated with respect to the available forest


and hence occupancy has remained constant in the area. Some areas between Katarniaghat and Suhelwa Wildlife Sanctuary have seen presence of leopard in 2022 where leopards were not reported in 2018 (Fig. II.1). Leopard occupied area in the landscape has remained consistent over the two monitoring cycles and only some parts of Nainital in Uttarakhand show loss in leopard occupancy in 2022 (Fig. II.1). Dehradun, Kalsi and Narendranagar Forest Division were sampled for the first time in 2022.

Leopard density was computed from 26 camera trapped sites within this landscape (Table II.1). A total of 10,564 leopard photo-captures were obtained from which 943 individuals were identified. Leopard sign encounter rate, prey density and human disturbance were used as covariates to model leopard density in a likelihood SECR framework. Model with leopard sign encounter and tiger sign best explained leopard density across the landscape. Total population of leopard within the sampled forested landscape of Shivalik-Gangetic plains was estimated at 1,109 (SE 134) as compared to 1,253 (SE 95) in 2018 and 929 (SE 75) in 2014.



©Trikanish Sharma

# Uttarakhand



**Figure II.3:**


*Spatially explicit leopard density (individuals /100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Uttarakhand, 2022.*

A total of 15 sites were camera trapped in Uttarakhand (Table II.1) that yielded 5,358 photo-captures of 588 leopard individuals. Like mentioned earlier, we only report leopard numbers from potential tiger habitat of the state. Leopard population of the state was 652 (SE 77) as compared to 839 (SE 48) in 2018. Leopard numbers remains stable in Rajaji and Corbett Tiger Reserve but density has not changed significantly since 2018 (Jhala *et al.*, 2021). Ramnagar Forest Division has recorded significant decline in leopard population possibly owing to increased tiger density (Qureshi *et al.* 2023). Three new sites, Kalsi, Dehradun and Narendranagar Forest Divisions were sampled for the first time. Dehradun Forest Division which was camera trapped for the first time had high leopard density and this is also the place where frequent movement of leopard is reported in the habitations. Terai West has seen significant increase in both tiger and leopard population. With frequent tiger attacks on people reported in the Terai (area south and east of Corbett Tiger Reserve) and a high density of leopard in the area as well, there is an urgent need to sensitise people about movement in forest. Both Terai West and Terai East are connected to Corbett and Pilibhit Tiger Reserve respectively and have the potential to be declared as Tiger

Reserves, this will assist in bringing much needed resources for better management.

Human leopard conflict remains cause of major concern in the state. With tiger and leopard occupying most of the forested areas, the state has to deal with conflicts on a regular basis. Leopard conflict is not limited to rural areas anymore and has spread to urban areas (like the recent leopard attacks in the outskirts of Dehradun city). Most of the studies on leopard conflict in the state have been carried out above 2000 msl which is beyond the study area for this report. But the conflict is no more limited to higher reaches of the state as evident from the recent leopard attack in township of Raipur in Dehradun in January 2024. In the past 5 years there have been nearly 2000 human-animal conflict (that includes injuries caused to humans and death), of this around 570 were attributed to leopard (Uttarakhand Forest Department 2024). In the state of Uttarakhand 30% of all the human death and injury cases by wildlife were caused by leopard (Uttarakhand Forest Department, 2024)

# Uttar Pradesh



**Figure II.4:** Spatially explicit leopard density (individuals /100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Uttar Pradesh, 2022.

A total of 9 sites were camera trapped in Uttar Pradesh (Table II.1) that yielded 4,282 photo-captures of 317 leopard individuals. Leopard population in the state was estimated at 371 (SE 54). Even though overall leopard population for the state has increased as compared to 2018 which was 316 (SE 39), (Jhala *et al.*, 2021), leopard population has shown significant decline in Dudhwa National Park and remained constant in Pilibhit and Suhelwa (Table II.1). There has been a significant increase in leopard population in Katarniaghat and Kishanpur WLS since 2018. Leopard density in Rani- pur has also gone up but that is due to better sampling effort in 2022 as compared to 2018. This increase in leopard population in Katarniaghat and Kishanpur are a worrying trend with respect to human-animal conflict in the state. Uttar Pradesh is already battling increasing number of tigers using multiple use matrix around its Protected Areas and with 47% of the leopard population present outside Protected Areas, the conflict intensifies.

Uttar Pradesh's human-leopard conflict problem is an outcome of its protected area's unique geography where most

of the Protected Areas are less than 10 km wide for much of their extent, with Katarniaghat just 4-6 km wide. The matrix surrounding these areas in all directions comprises of extensive agricultural land (sugarcane, wheat and paddy cultivation) and numerous villages with high human densities. Farmlands are essentially an extension of wildlife habitats for many mammals, especially when the crops are tall. In case of Katarniaghat, the farmlands to the north of the sanctuary in Nepal are fallow for several months of the year and there is little sugarcane cultivation. Many villages are at the forest edge. Thus, human reach within the forest is high, both because of these settlements and because the forests are accessed by residents of matrix villages. Local dependence on forest resources (primarily firewood and fodder) is also high, and large number of cattle are grazed in some grasslands within Katarniaghat. Wild prey is relatively low in the division (Chanchani *et al.*, 2014). For these reasons, the likelihood of humans encountering leopards both within the division and around is high. Given Protected Areas narrow width and high human pressure, it is remarkable that the areas still harbour populations of large




carnivores and herbivores – which can be attributed both to conservation measures and protection, and equally to the willingness of communities to coexist with wildlife in their neighbourhoods. However, the long-term persistence of these species rests at least in part on more effective conflict management, principally to significantly reduce encounters between humans and leopards. Bista *et al.*, (2019) study in Katarniaghat notes that unlike attacks by tigers which typically occur in areas with dense cover, 38% of the conflict with leopards occurred when the victim was either inside or adjacent to a home. Another 40% of the conflict was recorded in agricultural fields, and 11% of attacks were on people who were defecating in farmlands (a common practice in rural areas as homes generally do not have toilets or their use is minimal). The remaining cases were reported from forests, along village roads and other places. Most of these rescue operations of leopards are carried-out during

the monsoon and winter seasons as the cover availability increases in sugarcane farmlands and as harvesting period approaches. It is further evident that female leopards give birth to their litters in sugarcane fields during the winter season and use these farmlands to raise their cubs (Bista *et al.*, 2019).

Local forest department has been sensitising the local community with awareness programs. These program are aimed at spreading awareness amongst locals by sending messages on the phone to alert the user and by providing solar street lights to increase visibility in the conflict prone villages. Awareness building program is an effective tool to reduce conflict by bringing in a behavioural change in people.

## Bihar



**Figure II.5:**

*Spatially explicit leopard density (individuals /100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Bihar, 2022.*

Valmiki Tiger Reserve was the only camera trapped site in Bihar, where 924 leopard photo-captures yielded 78 leopard individuals. Leopard population in the state is 86 (SE 3)

which was 98 (SE 8) in 2018 (Jhala *et al.*, 2021). The leopard population needs to be monitored, as Valmiki have limited space and large tiger and leopard population will contribute to conflict with humans.

# Conservation Implications

Shivalik Hills and Gangetic Plains landscape recorded decline in leopard population and the five Tiger Reserves in this landscape had leopard density varying from 2.12 (SE 0.55) leopards per 100 km<sup>2</sup> in Dudhwa National Park to 14.07 (SE 1.57) leopards per 100 km<sup>2</sup> in Katarniaghat Wildlife Sanctuary. Detection corrected sex ratio was female biased in all the Tiger Reserves except Pilibhit. Uttarakhand

harbours major portion of the leopard population in the landscape, and the numbers will be more given that the hills of the state were not sampled for leopards. All states in this landscape are facing human-animal conflict with respect to tiger and leopard, this conflict will intensify in future and the state needs to take pro-active measures to avoid loss of life and property in such cases. With 65% of leopard population in the state present outside Protected Areas, training of forest staff in the forest divisions and territorial forests is first step towards managing the conflict. Timely compensations and awareness amongst local communities will also help alleviate the problem.



©Vinay Venugopal

**Table II.1:** Sampling details and leopard density parameter estimates using spatially explicit capture mark recapture analysis in a likelihood framework for sites in Shivalik Hills and Gangetic Plains Landscape, 2022.

State	Site	Model Space (km <sup>2</sup> )	Camera traps	Mt+1	Best Fit Model	D <sub>seccr</sub> /100 km <sup>2</sup> (SE)	g <sub>0</sub> Female (SE)	g <sub>0</sub> Male (SE)	σ Female (SE) (km)	σ Male (SE) (km)	P <sub>mix</sub> (SE)
Uttarakhand	Dehradun	1643	266	82	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	9.58 (1.08)	0.05(0.005)	0.06(0.01)	1.39(0.06)	2.10(0.10)	0.71:0.29(0.06)
Uttarakhand	Kalsi	756	151	25	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(-)	5.80(1.24)	0.03(0.005)		1.31(0.11)	2.82(0.26)	0.70:0.30(0.10)
Uttarakhand	Narendranagar	NA	11	5	NA	NA	NA	NA	NA	NA	NA
Uttarakhand	Rajaji	1487	426	151	-(, P <sub>mix</sub> -(sex), g <sub>0</sub> -(	12.32(1.01)	0.04(0.003)		1.31(0.03)		0.35:0.65(0.05)
Uttarakhand	Lansdowne	1389	181	54	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	8.78(1.33)	0.09(0.01)	0.09(0.01)	0.70(0.04)	2.54(0.15)	0.85:0.15(0.04)
Uttarakhand	Corbett	2017	514	113	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	7.46(0.70)	0.01(0.001)	0.03(0.002)	1.83(0.07)	1.62(0.04)	0.61:0.39(0.05)
Uttarakhand	Terai West	624	88	32	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	9.54(1.71)	0.12(0.03)	0.10(0.01)	0.82(0.08)	1.58(0.09)	0.47:0.53(0.10)
Uttarakhand	Almora	NA	13	1	NA	NA	NA	NA	NA	NA	NA
Uttarakhand	Ramnagar	NA	100	14	NA	NA	NA	NA	NA	NA	NA
Uttarakhand	Haldwani	1114	149	43	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	8.20(1.26)	0.09(0.01)	0.05(0.01)	0.91(0.04)	1.99(0.10)	0.69:0.31(0.07)
Uttarakhand	Terai Central	NA	15	1	NA	NA	NA	NA	NA	NA	NA
Uttarakhand	Champawat	NA	32	12	NA	NA	NA	NA	NA	NA	NA
Uttarakhand	Nainital	642	40	16	-(, P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	7.83(2.12)	0.28(0.04)		0.72(0.06)		0.36:0.64(0.15)
Uttarakhand	Terai East	1225	178	55	-(, P <sub>mix</sub> -(sex), g <sub>0</sub> -(	6.82(0.93)	0.02(0.002)		2.36(0.10)		0.45:0.55(0.08)
Uttarakhand	Amangarh	NA	45	8	NA	NA	NA	NA	NA	NA	NA
Uttarakhand	Dudhwa	1196	296	17	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	2.12(0.55)	0.01(0.01)	0.02(0.003)	1.29(0.33)	5.09(0.24)	0.56:0.44(0.14)
Uttarakhand	Katarniaghat	669	251	81	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	14.07(1.57)	0.004(0.0005)	0.01(0.001)	3.41(0.17)	3.90(0.16)	0.62:0.38(0.05)
Uttarakhand	Kishenpur	1127	167	25	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	3.01(0.71)	0.08(0.02)	0.04(0.01)	0.92(0.07)	6.45(0.28)	0.88:0.12(0.06)
Uttarakhand	South Kheri	NA	22	3	NA	NA	NA	NA	NA	NA	NA
Uttarakhand	Pilibhit	1218	308	29	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	2.92(0.55)	0.02(0.003)	0.03(0.002)	3.86(0.26)	4.18(0.12)	0.31:0.69(0.09)
Uttarakhand	Sohagbarwa	500	80	53	-(, P <sub>mix</sub> -(sex), g <sub>0</sub> -(	13.12(1.81)	0.12(0.01)		1.32(0.04)		0.55:0.45(0.08)
Uttarakhand	Suhelwa	668	75	40	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	7.91(1.29)	0.17(0.03)	0.11(0.02)	0.90(0.07)	2.10(0.13)	0.67:0.33(0.08)
Uttarakhand	Ranipur	510	99	61	-(, P <sub>mix</sub> -(sex), g <sub>0</sub> -(	16.93(2.21)	0.03(0.003)		1.64 (0.07)		0.52:0.48(0.070)
Bihar	Valmiki	1097	429	78	-(sex), P <sub>mix</sub> -(sex), g <sub>0</sub> -(sex)	7.79(0.89)	0.06(0.01)	0.04(0.003)	1.24(0.04)	2.80(0.08)	0.66:0.34(0.05)

SE: Standard error, D<sub>SECCR</sub>: Density estimate from Maximum Likelihood based spatially explicit capture recapture, σ : Spatial scale of detection function, g<sub>0</sub>: Magnitude (intercept) of detection function, P<sub>mix</sub>: Detection corrected estimate of proportion of female and male



# III: Central India and Eastern Ghats Landscape

Ujjwal Kumar, Rutu J. Prajapati, Vishnupriya Kolipakam, Jayanta Kumar Bora, Omkar Nar, Ayan Sadhu, Juri Roy, Shravana Goswami, Ashish Prasad, Swati Saini, Dhruv Jain, Krishna Mishra, Kainat Latafat, Manish A. Singanjude, Anup Kumar Pradhan, Gaurav A. Shinde, Anshuman Gogoi, Hemant Kamdi, Satya.P. Yadav, Yadvendradev V. Jhala and Qamar Qureshi



©Subharanjan Sen


## Central India and Eastern Ghats


The Central Indian and Eastern Ghats landscape encompasses the semi-arid zone of Rajasthan, deciduous and semi-evergreen forests forming a continuous expanse across the Deccan plateau, which includes the states of Maharashtra, Madhya Pradesh, Chhattisgarh, Jharkhand, and Odisha. It comprises sections of the Eastern Ghats, extending into Telangana, Andhra Pradesh, and Odisha. For administrative continuity and to avoid dividing the state of Maharashtra, the Sahyadri Hills of Maharashtra in the northern Western Ghats are included in this landscape. Conversely, certain portions of the Eastern Ghats in Tamil Nadu and Karnataka are excluded from this chapter, as they are addressed in the context of the Western Ghats landscape chapter. It is important to note that the leopard population in Rajasthan is specifically reported for tiger-occupied habitats within the Ranthambore, Sariska, Mukundara Hills, Ramgarh Visdhari Tiger Reserves and three Wildlife Sanctuaries (Kumbhalgarh, Shergarh and Bhainsrodgarh).

The Central Indian and Eastern Ghats landscape harbours the largest population of leopards fostered by diverse array of habitats ranging from semi-arid to dry and moist mixed deciduous forests and a rich prey base (Jhala *et al.*, 2021). The country wide genetic assessment of the leopard population suggests the Central Indian leopard population also has the highest genetic diversity (Jhala *et al.*, 2021).

The leopard population in this landscape is growing, owing much of its success to the protective measures under the umbrella of tiger conservation. The Tiger Reserves within this landscape also emerge as the major source sites for leopard populations. However, the expanding tiger population is not without consequences, as it exerts regulatory pressure on leopards. A recent study in Kanha revealed adverse effects of tigers on leopard populations (Kumar *et al.*, 2019). It is also evident in high leopard abundance estimates in the Tiger Reserves in the region such as Panna and Sariska where tiger abundance is low (Jhala *et al.*, 2019).

Figure III.1 Change in leopard distribution from 2018 to 2022 in the Central India and Eastern Ghats landscape





**Figure III.2** Spatially explicit leopard density modelled from camera trap based capture-mark-recapture and covariates of leopard sign, prey abundance and human disturbance within the Central India and Eastern Ghats landscape, 2022.

# Leopard Occupancy, Population Extent and Abundance

Leopards exhibited a notable increase in occupancy within surveyed cells (100 km<sup>2</sup>) compared to the cells in the previous estimation cycle of 2018 (Figure III.1). Total 8652 cells (100 km<sup>2</sup>) were sampled in 2022, out of which leopard presence was recorded in 30.87% (2,671) cells. Leopard presence was consistent in 1,751 cells in 2018 & 2022; leopards were not detected in 455 of previously occupied cells, whereas leopard presence was detected in 833 previously unoccupied cells in 2018 (Fig. III.1). Major gain in occupancy was reported from Madhya Pradesh & Rajasthan (Due to additional sampling areas).

The central Indian landscape, encompassing a total area of 1,10,653 km<sup>2</sup>, exhibits significant leopard occupancy distributed across four distinct patches. These patches are identified as follows:

- (a) The central block spans the entire states of Madhya Pradesh, Chhattisgarh, Jharkhand, Odisha, Maharashtra, and Northern Telangana.
- (b) The southern block extends from the Amrabad Tiger Reserve and Nagarjunsagar Srisailem Tiger Reserve to the Sri Venkateshwara Wildlife Sanctuary.
- (c) The western block includes the western ghats of Maharashtra (Sahyadri hills) and the surrounding areas of the Deccan Plateau.
- (d) The northern block encompasses Sariska, Ranthambore, and Mukundara Tiger Reserves, as well as the northern part of Madhya Pradesh, consisting of the Kuno-Palpur National Park, Madhav National Park, and Sheopur forests.

Leopard presence has been identified across all Protected Areas (Protected Areas) and major forest corridors within the central Indian landscape. Madhya Pradesh exhibits the highest leopard occupancy in this region, followed by Maharashtra. The comprehensive understanding of leopard distribution in these patches contributes to the conservation efforts and ecological management of leopard in Central India.

Leopard densities were computed using data from 67 camera-trapped sites in this region. A total of 51,598 photographs were obtained that yielded 3,666 unique individuals. Over 200 individual leopards were captured on camera in Panna, Nagarjunsagar Srisailem, and Satpura Tiger Reserves (Table III.1). The total leopard population in the surveyed forest of this area was estimated at 8,820 (SE 600). About 68% leopard population are outside to the Protected Area (PAs). However, high leopard densities were mainly observed in Protected Areas (PAs) and the major forest corridors connecting Protected Areas in this region (Figure III.2). The leopard population has increased in Madhya Pradesh, Maharashtra, and Andhra Pradesh, while a decline has been noted in Odisha, Chhattisgarh, and Telangana. It's noteworthy that the trend is similar to the decline in the tiger population (Qureshi *et al.*, 2023). This suggests common threats, such as poaching of prey for bush meat, targeted poaching for tiger and leopard skins and body parts (Raza *et al.*, 2012), and habitat loss due to mining and other human activities. Additionally, road accidents are a significant cause of leopard fatalities.


To address these issues, states should consider supplementing prey in low-density areas, implementing effective protection strategies using M-STrIPES, and providing wildlife management training for forest staff. Ensuring wildlife passage measures on roads passing through leopard habitats is also crucial.



©Vinay Venugopal



# Andhra Pradesh



**Figure III.3:**

*Spatially explicit leopard density (individuals/100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Andhra Pradesh, 2022*


Andhra Pradesh has a forest cover of 29,784 km<sup>2</sup>, making up 18.28 % of the state's total area (Indian State of the Forest Report 2021). Andhra Pradesh's diverse topography, ranging from the hills of Eastern Ghats and Nallamala to the shores of the Bay of Bengal, supports a rich variety of ecosystems. Eco-geographically the leopard population in Andhra Pradesh categorized into two regions (Figure III.3): (i) Rayalaseema: This region encompasses the forests of Nagarjunasagar Srisaillam Tiger Reserve (NSTR), Sri Venkateshwara National Park, Sri Penusila Narasimha Wildlife Sanctuary and the territorial areas of Kadapa Forest Divisions. (ii) Godawari basin: Leopard signs were not reported in this region, except for Papikonda National Park, situated in the Papi hills with its forested habitat extending into Odisha.

Leopard occupancy has been reported to have declined in the southern region of Papikonda National Park. However, there were mixed occupancy changes in the Rayalaseema region (Figure III.1). Camera trapping was carried out at Papikonda National Park and Nagarjuna Srisaillam Tiger Reserve, along with nearby sanctuaries including Sri Lankamalleswara, Sri Penusila Narasimha Wildlife Sanctuaries, Sri Venkateshwara

National Park, and the territorial areas of Kadapa Forest Divisions. A total of four sites were camera trapped in Andhra Pradesh that yielded of 310 unique leopard individuals. The total estimated population of Andhra Pradesh is 569 (SE 41) leopards in 2022 showed an increase compared to previous estimate 492 (SE 31) of countrywide monitoring of 2018 (Jhala *et al.*, 2021). Majority of the leopard population (69%) is concentrated in the Protected Areas and 31% population is in the territorial forests (Figure III.3). The leopard density from NSTR is not significantly different from previous estimate of 2018 (Jhala *et al.*, 2021), however more than 250 Adult leopards were photo-captured from NSTR alone (Table III.1), the largest single population.

A significant threat to the leopard population in Andhra Pradesh is human-leopard conflict. There have been reports of leopard attacks on devotees near the Tirumala hills, where the renowned Tirupati temple is situated. Other threats such as road kills by speeding vehicle on the road passing through leopard habitat are major concern in Andhra Pradesh. Appropriate mitigation measures should be adopted on the roads for the passageways of leopards.

# Chhattisgarh



**Figure III.4:**

*Spatially explicit leopard density (individuals/100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Chhattisgarh, 2022*


Chhattisgarh has a forest cover of 55,717 km<sup>2</sup> constituting 41.21% of the state's total area (Indian State of the Forest Report 2021). Although leopard distribution is interconnected within Chhattisgarh, the population and occupancy can be categorized into three blocks based on the connectivity and eco-geographic zones. (i) Northern Chhattisgarh, comprising Guru Ghasidas National Park-Tamor Pingla Wildlife Sanctuary, connected to Achanakmar Tiger Reserve through the forests of Madhya Pradesh. The area mostly falls under Northern hills and part of Chhattisgarh plains. Achanakmar Tiger Reserve holds the highest leopard population within the state's protected area network. (ii) Central Chhattisgarh, including Udanti Sitanadi Tiger Reserve and adjacent forest divisions along with Gomardha Wildlife Sanctuaries.

The region is located primarily within the Chhattisgarh plains and includes a portion of the Bastar hills. Leopard occupancy in these areas has declined since the previous nationwide monitoring exercise in 2018 (Fig. III.1). (iii) Southern Chhattisgarh, encompassing Indravati Tiger Re-

serve and the territorial forests of the Bastar region, connected to Gadchiroli in Maharashtra and the Kawal region of Telangana. The area mostly falls into Bastar plateau region of Chhattisgarh. The total estimated population of Chhattisgarh is 722 (SE 45) leopards in 2022, showing a decline compared to the previous estimate of 852 (SE 39) in 2018 (Jhala *et al.*, 2021). A total of two sites were camera trapped in Chhattisgarh that yielded 96 unique leopard individuals. Majority of the leopard population (75%) are outside PA network and concentrated in the territorial forests (Figure III.4).

The leopard density at Achanakmar Tiger Reserve was 7.19 (SE 0.89) per 100 km<sup>2</sup> which is not significantly different from previous estimate of 2018 (Jhala *et al.*, 2021) (Table III.1), however Udanti Sitanadi Tiger Reserve showed a decline (Table III.1). The leopard population and occupancy both are on declining trend in Chhattisgarh. The major threats are poaching (Rana & Kumar 2023), habitat loss, and prey depletion.

# Jharkhand



**Figure III.5:**

*Spatially explicit leopard density (individuals/100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Jharkhand, 2022.*


Jharkhand has a forest cover of 23,721 km<sup>2</sup> constituting 29.76 % of the state's total area (Indian State of the Forest Report 2021). The leopard population in Jharkhand is facing a worsening situation, with current occupancy only reported in Palamau Tiger Reserve and its adjacent forest division. Leopard distribution in Jharkhand was not uniform across the state, and their presence were commonly reported in Protected Areas such as Dalma and Hazaribagh Wildlife Sanctuaries besides Palamau Tiger Reserve. These regions provide suitable ecosystems for leopards, characterized by a mix of deciduous and tropical forests.

However, during the estimation of 2022, leopard occupancy from the other Protected Areas of Jharkhand were not reported (Figure III.1). The total estimated population of

Jharkhand is 51 (SE 10) leopards in 2022 (Figure III.5) which is stable in comparison to countrywide monitoring of 2018 (Jhala *et al.*, 2021). Only one site, i.e. Palamau Tiger Reserve was camera trapped in Jharkhand that yielded 23 unique leopard individuals. The leopard density at Palamau Tiger Reserve was reported to be 3.06/100 km<sup>2</sup> (Table III.1).

The state need to control prey poaching for bush meat consumption. There is a need to prioritize efforts to enhance the skill set of its forest staff through comprehensive capacity-building programs and to strengthen protection measures in National Parks and Sanctuaries. Furthermore, innovative methods which involves providing safe passage should be implemented to mitigate the impact of mining operations near forested areas.

# Madhya Pradesh



**Figure III.6:**


*Spatially explicit leopard density (individuals/100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Madhya Pradesh, 2022.*

Madhya Pradesh has a forest cover of 77,493 km<sup>2</sup> constituting 25.14 % of the state's total area (Indian State of the Forest Report 2021). Madhya Pradesh features diverse eco-geographic characteristics, encompassing parts of the semi-arid zone, central highlands, and plateau biotic provinces within the Deccan Peninsula biogeographic zones of India. Leopard has colonised in the Nauradehi Wildlife Sanctuary and northern part of Rani Duragawati Wildlife Sanctuary both are now part of the newly created Veerangana Durgawati Tiger Reserve.

Leopard occupancy has increased in the Chambal and Bundelkhand regions of Madhya Pradesh, with a marginal decline reported from the Chhindwara forests. Madhya Pradesh holds the largest leopard population in the country 3,907 (SE 215), (Table I.3) which is a significant increase from the previous estimation of 3,421 (SE 150) in 2018. Camera trapping at a total of 27 sites in Madhya Pradesh yielded images of 1692 unique leopard individuals (common individuals between sites counted only once). The majority of the leopard population (74%) exists outside

the Protected Area network, residing in territorial forest divisions or the multiple-use (buffer zone) forests of Tiger Reserves (Figure III.6). Panna Tiger Reserve and Kuno National Park exhibit some of the highest leopard densities in India (Table III.1). Other Tiger Reserves, including Satpura, Bandhavgarh, Pench, and Kanha, also feature high leopard densities, showcasing their coexistence with tigers. Obedullaganj (Ratapani Wildlife Sanctuary) and Balaghat forest divisions have reported a significant increase in leopard densities compared to the nationwide estimates of 2018 (Table III.1) (Jhala *et al.*, 2021). We believe due to data loss of leopard photos in 2018, the density estimates of Obedullaganj & Kuno were lower and should not be used for population trend with the 2022 estimates. While leopard populations thrive in Madhya Pradesh, ongoing effective vigilance is essential to address both targeted and unintentional poaching. Additionally, there is a need for the effective implementation of mitigation measures concerning linear infrastructure development projects to ensure the continued well-being of leopard populations in the region.

# Maharashtra



**Figure III.7:**


*Spatially explicit leopard density (individuals/100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Maharashtra, 2022.*

Maharashtra has a forest cover of 50,798 km<sup>2</sup> constituting 16.51 % of the state's total area (Indian State of the Forest Report 2021). Maharashtra spans three bio-geographic zones: The Deccan Peninsula, Western Ghats, and West Coast. In the central plateau region, dry deciduous mixed and dry teak forests are predominant, while the Western Ghats region is characterized by moist deciduous and evergreen forests. Leopards are distributed across Maharashtra's forested landscape, with limited presence was recorded in the large forest patches of Sahyadri region close to west coast (Figure III.1).

Maharashtra holds the second largest leopard population in the country 1,985 (SE 122), which is a significant increase from the previous estimation of 1,690 (SE 99) in 2018. Major population (75%) are outside the PA network (Figure III.7). Camera trapping at a total of 16 sites in Maharashtra yielded images of 904 unique leopard individuals (common individuals between sites counted only once). Based on the population connectivity, Leopard population in Maharashtra are classified into three sub population (i) Vidarbha : This region includes most of the Tiger Reserves,

such as Bor, Tadoba-Andhari, Nawegaon-Nagzira, Pench, and Melghat. It also encompasses sanctuaries like Painganga, Tipeswar, Umred-Pauni-Karhandla and large forest patches in territorial forest divisions such as Chandrapur, Central Chanda, Wardha, Yavatmal, and Gadchiroli. The leopard densities within this region is moderate to high. Leopard density has increased in the Chandrapur & Bramhapuri territorial divisions, Melghat and Tadoba Tiger Reserves since previous countrywide estimates of 2018 (Table III.1) (Jhala *et al.*, 2021). Tiger and leopard densities both are occurring at same scale within this region (Qureshi *et al.*, 2023). (ii) Sahyadri – Nasik: This area encompasses the Sahyadri Tiger Reserve, Sanjay Gandhi National Park, Radhanagri Wildlife Sanctuary, and the territorial forests of Sindhudurg, Kolhapur, Sawantwadi up to Nashik divisions. The leopard density of Sahyadri Tiger Reserve has increased from the previous countrywide estimate of 2018 (Table III.1) (Jhala *et al.*, 2021). We believe due to data loss of leopard photos in 2018, the density estimates of Sahyadri & Chandrapur were lower and should not be used for population trend with the 2022 estimates. Major threat for leopard in Maharashtra are conflict with humans and targeted poaching for body parts.

# Odisha



**Figure III.8:**

*Spatially explicit leopard density (individuals/100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Odisha, 2022*


Odisha has a forest cover of 52,156 km<sup>2</sup> constituting 33.50 % of the state's total area (Indian State of the Forest Report 2021). Odisha's major forest types include peninsular Sal and dry deciduous mixed forests. Leopard occupancy has been reported exclusively within Protected Areas (PAs) and their adjoining forest divisions. The leopard occupancy has declined from the previous occupied regions.

Leopard presence was not recorded from Nayagarh and Gumsur forest divisions in this cycle of countrywide monitoring. Leopard distribution is now majorly concentrated in the Simlipal & Satkosia Tiger Reserves, Hirakund, Kotagarh Khalasuni Wildlife Sanctuaries (Figure III.8). Camera trapping at total of three sites in Odisha yielded images

of 162 unique leopard individuals (common individuals between sites counted only once).

The leopard population in Odisha is estimated at 568 (SE 35), the population has significantly declined since last countrywide estimate of 760 (SE 33). However, the leopard densities from both the Tiger Reserves have increased significantly (Table III.1). This indicates the major efforts of recovery is restricted to Tiger Reserve. The state should also put efforts of effective patrolling, prey recovery and habitat management outside Tiger Reserve since major population of leopard (55%) are outside the Protected Area network (Figure III.8).

# Rajasthan



**Figure III.9:**

*Spatially explicit leopard density (individuals/100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Rajasthan, 2022.*


Rajasthan has a forest cover of 16,655 km<sup>2</sup> constituting 4.87 % of the state's total area (Indian State of the Forest Report 2021). The semi-arid zone of Rajasthan marks the westernmost extent of the central Indian landscape. The dry forests in this region are predominantly characterized by *Anogeisus pendula*, accompanied by associated species such as *Acacia*, *Butea*, *Lannea*, and others.

This terrain is significant for its distinctive topography and diverse flora, contributing to the ecological richness of the state. The sampling was restricted to Tiger Reserves and three wildlife sanctuaries (Kumbhalgarh, Bhainsrodgarh and Shergarh), sanctuaries were sampled first time in this countrywide monitoring exercise. These eight sites yielded 337 individual leopards. Leopard population of the state was

estimated at 721 (SE 112). Although leopard population in Rajasthan seems to be stable; however, direct comparisons with previous estimates are not possible due to the inclusion of new sampling areas in this nationwide estimation of 2022 (Figure III.9).

Sariska still has the highest leopard density of 21.43 (SE 1.91) amongst all the camera trapped sites in India. The density of leopard in Ranthambore Tiger Reserve has declined, however population remain stable (Table III.1). The primary issue facing leopards in this landscape is the loss and fragmentation of their habitat, primarily attributed to mining and developmental projects. To mitigate leopard-human conflict in the region, it is crucial to address the expansion of human land use and the escalating anthropogenic activities within leopard habitats.

# Telangana



**Figure III.10:**

*Spatially explicit leopard density (individuals/100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Telangana, 2022.*

Telangana has a forest cover of 21,214 km<sup>2</sup> constituting 18.93 % of the state's total area (Indian State of the Forest Report 2021). Telangana is situated in the Central Plateau biotic province of the Deccan Peninsula bio-geographic zone. The predominant forest types include dry deciduous mixed forests, dry deciduous scrub, and dry teak forests. The occupancy of leopard in Telangana was stable and mainly concentrated in the two Tiger Reserves (Kawal and Amrabad). Leopard has occupied new areas in northern Telangana, however leopard occupancy has declined from Kinnersani Wildlife Sanctuary (Fig. III.1).

Six sites were camera trapped in Telangana that yielded 146 individual leopards. Leopard population of the state was estimated at 297 (SE 20), which has declined from the previous countrywide estimate of 334 (SE 16) of 2018. Based on population connectivity leopard distribution can be divided into two population (i) Northern Telangana: This include Kawal Tiger Reserve, Kagaznagar forest division up to Pocharam Wildlife Sanctuary (Figure III.10). The leopard density in the Kawal was reported to be 1.19 (SE 0.29) per 100 km<sup>2</sup> (Table III.1). (ii) Southern Telangana: This includes Amrabad Tiger Reserve where Leopard density was 3.94 (SE 0.37) per 100 km<sup>2</sup> (Table III.1).





**Table III.1** Sampling details and leopard density parameter estimates using spatially explicit capture mark recapture analysis in a likelihood framework for sites in Central India and Eastern Ghats landscape, 2022

State	Site	Model space (km <sup>2</sup> )	Camera traps	M <sub>t+1</sub>	Best fit model	$\hat{D}$ secr/ 100 km <sup>2</sup> (SE)	g <sub>0</sub> Female (SE)	g <sub>0</sub> Male (SE)	σ Female (SE) Km	σ Male (SE) Km	Pmix (SE)
Andhra Pradesh	Nagarjunasagar Srisa- ilam TR	6211	783	254	σ <sup>~</sup> (sex), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (sex)	5.85 (0.38)	0.01 (0.001)	0.02 (0.001)	1.9 (0.05)	3.08 (0.07)	0.75:0.25 (0.03)
Andhra Pradesh	Narsimha WLS	1962	84	23	σ <sup>~</sup> (.), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	2.02 (0.47)	0.02 (0.004)	0.02 (0.004)	2.9 (0.33)	2.9 (0.33)	0.47:0.53 (0.13)
Andhra Pradesh	Papikonda	1220	103	20	σ <sup>~</sup> (sex), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	3.05 (0.8)	0.02 (0.01)	0.02 (0.01)	1.34 (0.18)	2.31 (0.31)	0.88:0.12 (0.07)
Andhra Pradesh	Venkateshwara NP	929	19	13	σ <sup>~</sup> (.), g0 <sup>~</sup> (.)	2.6 (0.09)	0.02 (0.011)	0.02 (0.011)	2.59 (0.48)	2.59 (0.48)	NA
Chhattis- garh	Achanakmar TR	1517	317	70	σ <sup>~</sup> (sex), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (sex)	7.19 (0.89)	0.02 (0.003)	0.028 (0.003)	1.52 (0.07)	2.83 (0.12)	0.79:0.21 (0.05)
Chhattis- garh	Udanti-Sitanadi TR	2684	142	26	σ <sup>~</sup> (sex), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	1.97 (0.44)	0.024 (0.005)	0.024 (0.005)	2.24 (0.28)	3.5 (0.4)	0.7:0.3 (0.11)
Jharkhand	Palamau TR	1664	323	23	σ <sup>~</sup> (.), g0 <sup>~</sup> (.)	3.06 (0.76)	0.003 (0.001)	0.003 (0.001)	3.18 (0.43)	3.18 (0.43)	NA
Madhya Pradesh	Balaghat	3278	495	134	σ <sup>~</sup> (sex), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	7.15 (0.66)	0.01 (0.001)	0.01 (0.001)	1.31 (0.05)	2.94 (0.1)	0.7:0.3 (0.04)
Madhya Pradesh	Bandhavgarh TR	2038	697	133	σ <sup>~</sup> (.), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	8.65 (0.76)	0.01 (0.001)	0.01 (0.001)	1.93 (0.06)	1.93 (0.06)	0.59:0.41 (0.05)
Madhya Pradesh	Barghat	1013	69	20	σ <sup>~</sup> (.), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	5.85 (2.1)	0.02 (0.01)	0.02 (0.01)	2.41 (0.56)	2.41 (0.56)	0.5:0.5 (0.14)
Madhya Pradesh	North Betul	710	107	22	σ <sup>~</sup> (sex), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (sex)	8.94 (3.15)	0.003 (0.003)	0.02 (0.004)	2.26 (1.25)	5.53 (0.49)	0.84:0.16 (0.08)
Madhya Pradesh	Bhopal (Samardha)	258	43	8	σ <sup>~</sup> (.), g0 <sup>~</sup> (.)	6.46 (2.71)	0.01 (0.003)	0.01 (0.003)	1.32 (0.27)	1.32 (0.27)	NA
Madhya Pradesh	Dewas	741	120	21	σ <sup>~</sup> (.), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	3.81 (0.87)	0.01 (0.003)	0.01 (0.003)	3.32 (0.34)	3.32 (0.34)	0.46:0.54 (0.14)
Madhya Pradesh	Kanha TR	2430	664	153	σ <sup>~</sup> (sex), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	8.63 (0.71)	0.02 (0.002)	0.02 (0.002)	1.7 (0.04)	2.88 (0.07)	0.72:0.28 (0.04)
Madhya Pradesh	Kuno WLD	1007	146	111	σ <sup>~</sup> (.), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	16.38 (1.59)	0.04 (0.003)	0.04 (0.003)	1.27 (0.04)	1.27 (0.04)	0.54:0.46 (0.05)
Madhya Pradesh	Lamta	1573	221	84	σ <sup>~</sup> (sex), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (sex)	9.29 (1.11)	0.02 (0.003)	0.03 (0.003)	1.33 (0.09)	2.86 (0.11)	0.67:0.33 (0.05)
Madhya Pradesh	Mandla	942	107	28	σ <sup>~</sup> (sex), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (sex)	5.26 (1.13)	0.02 (0.004)	0.03 (0.005)	1.63 (0.21)	3.1 (0.23)	0.73:0.27 (0.08)
Madhya Pradesh	Narmadapuram	1857	267	29	σ <sup>~</sup> (.), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	3.49 (0.8)	0.002 (0.001)	0.002 (0.001)	2.97 (0.41)	2.97 (0.41)	0.67:0.33 (0.1)
Madhya Pradesh	Nauradehi WLS	NA	114	1	NA	NA	NA	NA	NA	NA	NA
Madhya Pradesh	North Panna	457	112	40	σ <sup>~</sup> (.), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	11.8 (1.91)	0.06 (0.01)	0.06 (0.01)	1.07 (0.06)	1.07 (0.06)	0.54:0.46 (0.1)
Madhya Pradesh	Obedullaganj -Rata- pani WLS	2515	666	134	σ <sup>~</sup> (.), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	6.64 (0.58)	0.01 (0.001)	0.01 (0.001)	2.1 (0.06)	2.1 (0.06)	0.42:0.58 (0.05)
Madhya Pradesh	Panna TR	1977	574	240	σ <sup>~</sup> (sex), Pmix <sup>~</sup> (sex), g0 <sup>~</sup> (.)	16.09 (1.05)	0.01 (0.001)	0.01 (0.001)	1.41 (0.04)	2.19 (0.05)	0.56:0.44 (0.03)

State	Site	Model space (km <sup>2</sup> )	Camera traps	M <sub>t+1</sub>	Best fit model	$\hat{D}_{sec}/100\text{ km}^2$ (SE)	$\xi_0$ Female (SE)	$\xi_0$ Male (SE)	$\sigma$ Female (SE) Km	$\sigma$ Male (SE) Km	Pmix (SE)
Madhya Pradesh	Pench TR	1440	426	125	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	12.27 (1.11)	0.02 (0.001)		1.37 (0.04)	2.54 (0.08)	0.74±0.26 (0.04)
Madhya Pradesh	Rampur Bhatodi	608	39	11	$\sigma \sim (.), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	3 (1.12)	0.01 (0.004)		3.59 (0.67)		0.44±0.56 (0.17)
Madhya Pradesh	Sanjay Dubari TR	2504	573	104	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	6.17 (0.62)	0.013 (0.002)	0.009 (0.001)	1.88 (0.09)	3.67 (0.15)	0.70±0.30 (0.05)
Madhya Pradesh	Satna	504	88	25	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	10.35 (2.54)	0.02 (0.005)		0.97 (0.14)	2.65 (0.26)	0.71±0.29 (0.1)
Madhya Pradesh	Satpura TR	2510	830	210	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	10.23 (0.71)	0.03 (0.001)		1.49 (0.03)	2.27 (0.04)	0.65±0.35 (0.03)
Madhya Pradesh	Sehore	455	48	10	$\sigma \sim (.), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	3.53 (1.21)	0.04 (0.01)		1.75 (0.24)		0.33±0.67 (0.16)
Madhya Pradesh	Shahdol	1456	95	17	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	4.82 (1.86)	0.01 (0.003)		1.83 (0.45)	4.1 (0.79)	0.81±0.19 (0.11)
Madhya Pradesh	South Panna	1141	94	36	$\sigma \sim (.), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	6.09 (1.16)	0.03 (0.01)		1.63 (0.15)		0.39±0.61 (0.11)
Madhya Pradesh	South Seoni	937	96	18	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	6.4 (1.97)	0.01 (0.003)		1.77 (0.31)	3.9 (0.77)	0.87±0.13 (0.08)
Madhya Pradesh	Umaria	669	102	19	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	6.42 (1.66)	0.02 (0.004)		1.33 (0.17)	2.43 (0.33)	0.81±0.19 (0.09)
Madhya Pradesh	West Betul	751	100	16	$\sigma \sim (.), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	5.5 (2.14)	0.001 (0.001)	0.03 (0.01)		3.37 (0.34)	0.79±0.21 (0.1)
Maharashtra	Bor TR	880	206	34	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	5.14 (0.9)	0.03 (0.003)		1.5 (0.1)	2.36 (0.15)	0.53±0.47 (0.09)
Maharashtra	Bramhapuri	1931	417	106	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	8.16 (0.82)	0.015 (0.002)	0.023 (0.002)	1.47 (0.07)	3.43 (0.1)	0.73±0.27 (0.04)
Maharashtra	Central Chanda	1515	308	50	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	4.43 (0.64)	0.01 (0.001)	0.03 (0.002)	2.88 (0.16)	3.71 (0.13)	0.65±0.35 (0.07)
Maharashtra	Chandrapur	1223	223	69	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	8.01 (0.99)	0.03 (0.003)	0.04 (0.004)	1.51 (0.08)	2.8 (0.16)	0.75±0.25 (0.05)
Maharashtra	Dnyanganga	NA	82	44	NA	NA	NA	NA	NA	NA	NA
Maharashtra	Gadchiroli	4215	541	47	$\sigma \sim (.), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	1.8 (0.27)	0.005 (0.001)	0.015 (0.001)		5.07 (0.2)	0.62±0.38 (0.07)
Maharashtra	Karanja Sohul WLS	NA	8	2	NA	NA	NA	NA	NA	NA	NA
Maharashtra	Katepurna WLS	59	15	7	$\sigma \sim (.), g0 \sim (.)$	13.33 (5.33)	0.02 (0.01)		2.54 (0.78)		NA
Maharashtra	Melghat TR	3300	885	171	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	7.11 (0.55)	0.02 (0.002)	0.02 (0.001)	1.57 (0.06)	2.57 (0.06)	0.67±0.33 (0.04)
Maharashtra	Navegaon-Nagzira TR	1758	526	110	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	8.03 (0.77)	0.009 (0.001)	0.012 (0.001)	1.61 (0.06)	2.9 (0.1)	0.75±0.25 (0.04)
Maharashtra	Painganga WLS	430	42	11	$\sigma \sim (.), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	3.89 (1.3)	0.02 (0.005)			2.27 (0.4)	0.25±0.75 (0.15)
Maharashtra	Pandharkawada	1099	106	10	$\sigma \sim (.), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	1.15 (0.38)	0.01 (0.002)			7.68 (1.13)	0.43±0.57 (0.19)
Maharashtra	Pench TR	1172	302	64	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (.)$	8.79 (1.14)	0.02 (0.002)		1.45 (0.08)	2.74 (0.13)	0.7±0.3 (0.06)
Maharashtra	Sahyadri TR	2648	435	83	$\sigma \sim (.), g0 \sim (.)$	5.09 (0.59)	0.003 (0.0003)			3.25 (0.15)	NA

State	Site	Model space (km <sup>2</sup> )	Camera traps	M <sub>t+1</sub>	Best fit model	$\hat{D}$ secr/100 km <sup>2</sup> (SE)	g <sub>0</sub> Female (SE)	g <sub>0</sub> Male (SE)	$\sigma$ Female (SE) Km	$\sigma$ Male (SE) Km	Pmix (SE)
Maharashtra	Tadoba Andhari TR	1848	630	129	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\cdot)$	8.86 (0.78)	0.02 (0.001)	3.32 (0.07)	1.68 (0.04)	3.32 (0.07)	0.7:0.3 (0.04)
Maharashtra	Tipeswar WLS	NA	62	5	NA	NA	NA	NA	NA	NA	NA
Maharashtra	Umred Pauni Karhandla WLS	225	85	7	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g0 \sim (\cdot)$	3.38 (1.33)	0.03 (0.005)	3.43 (0.66)	3.43 (0.66)	3.43 (0.66)	0.4:0.6 (0.22)
Odisha	Kuldiha WLS	NA	16	0	NA	NA	NA	NA	NA	NA	NA
Odisha	Satkosia TR	1493	258	77	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g0 \sim (\cdot)$	7.41 (0.86)	0.02 (0.016)	2.02 (0.07)	2.02 (0.07)	2.02 (0.07)	0.5:0.5 (0.07)
Odisha	Similipal TR	2330	459	85	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	5.67 (0.64)	0.01 (0.001)	0.03 (0.002)	1.81 (0.08)	3.31 (0.12)	0.8:0.2 (0.04)
Rajasthan	Bhainsrodgarh WLS	365	29	10	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g0 \sim (\cdot)$	8.93 (3.82)	0.035 (0.015)	1.03 (0.21)	1.03 (0.21)	1.03 (0.21)	0.6:0.4 (0.22)
Rajasthan	Kailadevi WLS (RTR II)	NA	14	5	NA	NA	NA	NA	NA	NA	NA
Rajasthan	Kumbhalgarh WLS	533	52	55	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\cdot)$	19.20 (2.68)	0.12 (0.012)	1.75 (0.098)	0.82 (0.04)	1.75 (0.098)	0.72:0.28 (0.06)
Rajasthan	Mukundara Hills TR	2661	169	41	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\cdot)$	4.87 (0.088)	0.0116 (0.003)	2.83 (0.33)	1.55 (0.016)	2.83 (0.33)	0.73:0.27 (0.09)
Rajasthan	Ramgarh Visdhari TR	811	74	19	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g0 \sim (\cdot)$	3.11 (0.75)	0.032 (0.006)	2.58 (0.19)	2.58 (0.19)	2.58 (0.19)	0.33:0.67 (0.16)
Rajasthan	Ranthambore TR I	2034	158	66	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\cdot)$	12.97 (1.72)	0.02 (0.003)	1.51 (0.078)	0.79 (0.057)	1.51 (0.078)	0.62:0.38 (0.07)
Rajasthan	Sariska TR	1262	158	138	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	21.43 (1.91)	0.09 (0.008)	6.57 (1.95)	0.91 (0.029)	6.57 (1.95)	0.95:0.05 (0.04)
Rajasthan	Shergarh WLS	NA	31	3	NA	NA	NA	NA	NA	NA	NA
Telangana	Amrabad TR	4405	812	117	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g0 \sim (\text{sex})$	3.94 (0.37)	0.008 (0.001)	6.32 (0.22)	2.85 (0.1)	6.32 (0.22)	0.73:0.27 (0.04)
Telangana	Chennur	NA	151	0	NA	NA	NA	NA	NA	NA	NA
Telangana	Eturnagaram WLS	NA	96	0	NA	NA	NA	NA	NA	NA	NA
Telangana	Kagaznagar	1891	147	8	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g0 \sim (\cdot)$	0.52 (0.19)	0.02 (0.004)	7.36 (0.91)	7.36 (0.91)	7.36 (0.91)	0.33:0.67 (0.19)
Telangana	Kawal TR	2079	316	19	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g0 \sim (\cdot)$	1.19 (0.29)	0.01 (0.001)	5.81 (0.57)	5.81 (0.57)	5.81 (0.57)	0.69:0.31 (0.13)
Telangana	Kothagudem	NA	56	2	NA	NA	NA	NA	NA	NA	NA

SE: Standard error,  $\hat{D}^{\text{SECR}}$ : Density estimate from Maximum Likelihood based spatially explicit capture recapture,  $\sigma$ : Spatial scale of detection function,  $g0$ : Magnitude (intercept) of detection function, Pmix: Detection corrected estimate of proportion of female and male



# Conservation Implications


Leopards possess a broad dietary niche and has remarkable adaptability to a varying habitat condition, even in human dominated areas. A large (68%) population of leopard reside outside the Protected Areas in multiple use forests (Jhala *et al.*, 2021). This proximity to human habitation makes them more vulnerable to poaching and human leopard conflict. Among the large carnivores found in India, leopards are the most poached species (Mondol *et al.*, 2015). The Central Indian landscape stands out as a hotspot for leopard poaching, with a bias towards the poaching of males (Mondol *et al.*, 2015). Poaching of large carnivores may alter demography, behaviour of the target species also leaves cascading impact on the native biodiversity.

Beyond poaching, habitat fragmentation due to developmental activities such as the development of linear infrastructure and mining poses significant threats to leopards in this landscape. Rich in coal and mineral reserves, this region serves as a crucial link connecting major economic hubs through a network of roads and railways that intersect its biodiverse forest habitats. Over the last century, leopards in India have experienced a substantial human-induced population decline (Bhatt *et al.*, 2020).

Interestingly, this landscape experiences relatively fewer conflicts compared to hilly regions like Uttarakhand and Himachal Pradesh (Shivakumar *et al.*, 2023). The diverse habitat and abundant prey base in this landscape could be the reason of low conflict (Jhala *et al.*, 2015). However, as human populations encroach further into leopard habitats, incidents of human-leopard conflict are on the rise. In some instances, these conflicts escalate into violence, with communities retaliating against leopards that have preyed on their livestock or injures human. Maharashtra has emerged as the worst affected state, reporting 113 fatal attacks in the last seven years (Kulkarni, 2023).

While recent studies have accumulated substantial knowledge about leopard populations, habitat use, food habits, and human-leopard conflict, there remains a notable gap in understanding their behaviour in varying natural conditions. With the leopard population expected to increase in the future, a concerted effort toward long-term monitoring, coupled with a deeper understanding of their natural history and behaviour, is imperative to devise effective management strategies.





©Achintya Singh





# IV: Western Ghats Landscape

Ayan Sadhu, Ujjwal Kumar, Genie Murao, Vishnupriya Kolipakam, Swati Saini, Ashish Prasad, Kausik Banerjee, Prayas Auddy, Deepali Chatrath, Aritra Roy, Rajrajeshwar Thaker, Monika Saraswat, Vaishnavi Gusain, N.S Murali, Harini Venugopal, Rajendra Garawad, Amit Mallick, Satya P. Yadav, Yadvendradev V. Jhala and Qamar Qureshi



©Deeksha Natraj

## Western Ghats

Leopards are widely distributed in the southern states of Goa, Karnataka, Kerala and Tamil Nadu, from the semi-evergreen patches of Western Ghats to semi-arid landscapes of central and parts of northern Karnataka. Unlike tigers, leopards are often found in the human-dominated land uses in Western Ghats, having adjusted themselves in the mosaics of coffee estates, plantations, and forests (Athreya *et al.*, 2013, Gubbi *et al.*, 2017).

While in the forested landscapes, leopards majorly rely on wild prey species (Karanth and Sunquist 1995, Ramesh *et al.*, 2009), in the semi-urban set up, they can sustain themselves by consuming livestock, domestic dogs, and human subsidized food sources (Athreya *et al.*, 2014, Sidhu *et al.*, 2015). Such interfaces often result in conflicts with human (attack on human, livestock depredation) (Sidhu *et al.*, 2017). In recent years, numerous developmental projects, linear infrastructure and human encroachment caused habitat loss and fragmentation in the leopard distribution range (Gubbi *et al.*, 2017).

The leopard population in the landscape is almost continuous apart from a few natural breaks. The population in the northern part of Western Ghats (Kali-Sahyadri-Shi-

moga-Mollem cluster) is connected to Bhadra-Bhadra-vathi-Chikmagalur-Kudremukh cluster through the hill forests of Sharavathi-Someswara Wildlife Sanctuary (WLS). This landscape population further extends to Nilgiri cluster (Nagarhole, Wayanad, Bandipur, Mudumalai, BRT Hills, Sathyamangalam) through Brahmagiri-Pushpagiri-Talaka-veri Sanctuaries. Similar to tiger and elephant, the Nilgiri cluster harbors the single largest leopard population in India distributed amongst the states of Karnataka, Tamil Nadu, and Kerala, and serves as a link between Western Ghats and Eastern Ghats. In the north, the Nilgiri population further extends into the patchy forest-scrubland mosaics of Ramanagara, Bengaluru Urban, Bengaluru Rural, and Tumkur divisions through MM Hills, Cauvery Wildlife Sanctuary.

The central Karnataka leopard population is sparsely distributed into the forest-scrubland-agriculture mosaics of Ballari, Davangere divisions and connected through the stepping stone forest/refuge patches to Haveri, Dharwad and adjoining Divisions. Below the Silent Valley-Mukurthi forests (southern part of the Nilgiri cluster), the Palghat gap remains a geographical barrier and separates the Nilgiri cluster from the Anamalai-Parambikulam cluster. Further south, the semi-evergreen forests of Periyar-Srivilliputhur-Meghamalai (below Shencottah gap), Anamalai-Parambikulam in the north and the Kalakad Mun-

danthurai-Kanyakumari in the south forms important clusters of reasonable leopard population. The Kalakad Mundanthurai-Kanyakumari cluster forms the southernmost leopard population in India, distributed in the states of Tamil Nadu and Kerala. These two blocks south of the Palghat gap are connected through narrow linkages which are conducive to leopard movement between these two populations.

## Leopard Occupancy, Population Extent and Abundance

Leopards are one of the most versatile large carnivores in the world, known to occur in high densities in the prey rich undisturbed forests as well as managed to survive in the urban landscapes at the edge of forest patches. The forests of Western Ghats are some of the most productive forests in the country and harbor rich assemblages of predator and their prey (Jhala *et al.*, 2021).

As a part of the All India Leopard Estimation exercise (2022), a total of 1,558 cells (100 km<sup>2</sup>) were sampled using camera traps and foot sampling, out of which leopard was found to be present in 64% (1,001) cells. The camera trapping exercise resulted into 18,712 leopard photographs from which 2,052 individual leopards were identified. The spatially-explicit capture recapture estimate of leopard population in the Western Ghats was estimated at 3,596 (SE 482) leopards in the Western Ghats. Leopard presence in the landscape remains almost the same compared to 2018

(Fig. IV.1 & IV.2). Leopards occur in much lower densities in the scrubland-open forest mosaics of central Karnataka or the evergreen patches of southern Western Ghats (<1 leopard/100 km<sup>2</sup>) (Table IV.1). The latest All India Leopard Estimate exercise depicted, the Kali-Shimoga-Mollem cluster harbours a low to medium density leopard populations, with moderately high leopard density observed at the Kali Tiger Reserve (Fig. IV.2). Similar pattern can be observed in the Bhadra-Bhadravathi-Kudremukh cluster, where leopards mostly have a sporadic distribution in the landscape, and occur in high densities only within Bhadra TR (Fig. IV.2). In the more human-dominated areas of central Karnataka, leopards occur mostly at low densities. Leopards have been found in very high (12-13 leopards/100 km<sup>2</sup>) to moderately high densities in the Nilgiri cluster (Nagarhole-Wayanad-Bandipur-Mudumalai-BRT-Sathyamangalam TRs) (Fig. IV.2). The remaining forested habitats of the Nilgiri block also have moderate to high leopard densities, making this cluster a stronghold of leopard population. Its noteworthy to mention that the Nilgiri cluster also harbours the world's largest tiger population. The Anaimalai-Parambikulam cluster (south to Palghat gap) harbors moderate density leopard population, with leopards occurring at moderately high densities mostly in Parambikulam Tiger Reserve. Leopards occur at lower densities in the evergreen and semi-evergreen forests of Periyar-Srivilliputhur-Meghamalai landscape and KMTR-Kanyakumari landscape (Fig. IV.2).


The population estimate of the previous cycle (2018) estimated 3,387 (SE 142) leopards in the landscape (Jhala *et al.*, 2021). Within the consistently sampled area (sampled in 2018 and 2022), there is a marginal increase in the leopard population (133 leopards). Approximately 65% of this leopard population occurs outside the Protected Areas of the landscape.



©Vinay Venugopal


### Figure IV.1:


Change in leopard distribution from 2018 to 2022 in the Western Ghats Landscape.



**Figure IV.2:**

*Spatially explicit leopard density (individuals /100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates of leopard sign, prey and human disturbance for Western Ghats Landscape, 2022.*






**Figure IV.3:**

*Spatially explicit leopard density (individuals /100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Goa, 2022.*

Goa has a forest cover of approximately 1,800 km<sup>2</sup> which is contiguous with Kali Tiger Reserve in Karnataka. The state forms a linkage between the forests of the northern and central Western Ghats. For the first time as a part of the All India Tiger Estimation exercise the Goa Forest Department sampled 5 sites using camera traps that yielded 219 photo-captures of 30 individual leopards (Fig. IV.3). The leopard population of the state was estimated to be 77 (SE 13), from camera trap and covariate model, with approximately 26% of the population residing outside the Protected Areas of Goa. The continuous forest patches of Mhadei-Mollem harbor higher density of leopards compared to other parts of the state (Table IV.1, Fig. IV.3).

In Goa, there has been an increase in the cases of leopards straying into human habitations around the villages on the fringe of Cotigao Wildlife Sanctuary, that is, Loliem, Poinguinim, Cotigao, Gaondogrem, Sristhal, Agonda and Cola. Leopard mortalities or injuries due to snaring is not

uncommon as locals often put snares to capture wild pigs. Leopards also got killed by speeding trains and vehicles, an emerging threat to the wildlife of the state due to rapid development of unmitigated linear infrastructure. Leopards, sloth bears and gaurs often venture into cashew plantations which are in proximity to the forested landscapes, leading to increased instances of negative human-animal confrontations in residential areas. The recent increase in leopards straying into human habitations has raised concerns amongst the locals. It's important to investigate the state of the Goa's forests in terms of prey, habitat conditions, protection to gain insight into the situation. The Forest Department should initiate regular camera trapping for carnivores and monitoring of prey and their habitat in the forested landscapes of Goa apart from the All India Tiger Estimation. With appropriate management and conservation measures, the forests of Goa have immense potential for harboring sizeable large carnivore population.



**Figure IV.4:**

*Spatially explicit leopard density (individuals /100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Karnataka, 2022.*

The forests of Karnataka span over an area of 33,931 km<sup>2</sup> (Forest Survey of India 2017), encompassing a diverse landscape that features dry deciduous, moist deciduous, evergreen, semi-evergreen forests, scrub forests and rocky outcrops (Gubbi *et al.*, 2020). The state consists of three biogeographic zones, namely, the Coasts, Western Ghats and Eastern Plains (Deccan south) (Jhala *et al.*, 2021). The Karnataka Forest Department, under the ambit of All India Tiger Estimation, put enormous effort to sample 37 sites using camera traps yielded 9,351 photo-captures of 1,062 individual leopards (Fig. IV.4). The leopard population of the state remain stable with a pop-

ulation estimate of 1,879 leopards (SE 261) in 2022 compared to 2018 estimates of 1,783 leopards (SE 71) (Jhala *et al.*, 2021). The Nagarhole-Bandipur-BRT Hills Tiger Reserves harbor the largest leopard population in the state, while Bhadra Tiger Reserve holds the highest density of leopards (Table IV.1). Approximately 59% of the leopard population of Karnataka resides outside the Protected Areas, sporadically distributed in the territorial divisions of Tiger Reserves and the human-dominated areas of state, like Bengaluru rural, Dharwad, Davanagere, Hassan, Haveri, Koppa, Mangaluru, Ramanagara, Tumkur, Sagara, Sirsi and Virajpet. From 2018 to 2022, major-

# Karnataka

ity of the sampled sites in Karnataka showed stable population trends for leopards. Leopard densities remain stable in Bandipur, Belagavi, Bhadra, Bhadravathi, Cauvery, Kali, Nagarahole (Table IV.1). A significant increase in leopard population was observed in Bannerghatta National Park, where there has been a considerable increase in the spatial coverage of camera trapping. In BRT Hills, the leopard population had increased compared to 2018. In Bhadra Tiger Reserve and Kudremukh National Park, the number of individual leopard photo-captured increased to a great extent, over the cycles, but the densities have remained stable. The sampling area and spatial coverage of camera trapping increased significantly (Table IV.1). A sizable population of leopards in Karnataka inhabit areas in the proximity to human settlements, resulted in high numbers of negative interactions between human and leopards throughout the state (Gubbi *et al.*, 2017).


The conflict between humans and leopards, initially confined to mostly Ramanagara, Tumkuru, Mandya, Mysuru, and Hassan, however, gradually spread across to Ballari, Koppala, and Kolara. This escalation is attributed to extensive mining and widespread granite quarrying activities, leading to a significant reduction in the natural habitats of leopards (Reddy *et al.*, 2019). The adverse effects of dam constructions, the establishment of major roads through Protected Areas, soil erosion, mining operations, agricultural encroachment along forest boundaries, grazing of domestic animals, reduced rainfall, and climate change are likely to contribute to the diminishing wild spaces in the state.

In recent years, leopards have increasingly preyed on domestic animals within human habitats due to the decline in herbivore populations (Sidhu *et al.*, 2017), which has resulted in key species such as leopards and tigers finding themselves in conflicts with humans in the peripheral regions surrounding Protected Areas as well as territorial areas (Chauhan *et al.*, 2021). During the period from 2009 to 2016, the majority of conflict cases (~80%) were concentrated in the districts of Mysuru, Udupi, Hassan, Tumkuru, Ramanagara, Ballari, Koppala, and Mandya (Manjunatha *et al.*, 2023). Karnataka accounted for almost half of the reported wildlife conflict related deaths in the country during 2012-13 (Karanth *et al.*, 2013). From 2018 to 2023 there has been a gradual rise in the leopard conflict cases recorded in the state of Karnataka. In this time frame over 100 cases of human-leopard encounters were recorded in the state, with the highest number of incidents documented between November 2022 and January 2023 (Manjunatha *et al.*, 2023). Leopards were responsible for the death of around 516 domestic animals in Mandya district alone from 2019 to 2022 (Manjunatha *et al.*, 2023). There has been a recent surge in human fatalities and domestic animal killings attributed to leopard attacks, particularly in the taluks of Mysuru, Nanjangud, HD Kote, Sargur, T. Narsipur, Mandya, Pandavapura, Nagamangala, and the surrounding areas. Effective and timely management actions and mitigating the human-leopard conflict is the need of the hour in Karnataka in order to safeguard the leopard populations in the state.



©Gurinderjit Singh

# Tamil Nadu



**Figure IV.6:**

*Spatially explicit leopard density (individuals /100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Tamil Nadu, 2022.*

Tamil Nadu comprises three biogeographic zones, namely, Western Ghats, Deccan peninsula and Coastal zone. The state extends over both Western Ghats and Eastern Ghats, comprising the forested area of 21,509 km<sup>2</sup>. Nine out of 16 major forest types and 48 sub types of forest recognized by Champion and Seth (1968) occur in Tamil Nadu. A total of 14 sites were sampled using camera traps in Tamil Nadu that yielded 5,433 photo-captures of 736 individual leopards (Fig. IV.6). The leopard population of the state has significantly increased with an estimated 1,070 (SE 132) leopards in 2022, as compared to 868 (SE 40) leopards in 2018 (Jhala *et al.*, 2021) and 815 (SE 228) leopards in 2014 (Jhala *et al.*, 2015). A majority of the leopard population of Tamil Nadu, that is approximately 80%, reside outside the Protected Areas of the state, with high density leopard populations

mostly concentrated near the Nilgiri region (including Coimbatore, Erode, Gudalur and Nilgiri divisions) and in the Valparai area. Mukurthy, Kanyakumari and Kodaikanal exhibit consistently low densities of leopards.

From 2018 to 2022, the leopard populations of Erode, Gudalur, Mudumalai TR, Kanyakumari Wildlife Sanctuary, Kodaikanal Wildlife Sanctuary, Mukurthy National Park, Nellai Wildlife Sanctuary, Sathyamangalam and Srivilliputhur-Meghamalai Tiger Reserves have remained stable (Table IV.1). In case of Nilgiri and Meghamalai, there has been a significant increase in the sampling area and number of individual leopard photo-captured, however, the densities have remained stable. There has been a considerable increase in leopard density of Anamalai. On the other hand, KMTR




# Tamil Nadu


experienced a significant decline which may be a result of a decrease in spatial coverage of camera traps.

In Tamil Nadu, the coffee-tea estates and other commercial plantations surrounded by forests are frequently occupied by leopards and are major hubs for human-leopard conflicts (Sidhu *et al.*, 2017; Deivanayaki & Ezhilarasi 2019, Jhala *et al.*, 2021). Several cases have been reported from the forest

fragments of Valparai plateau and Anamalai Tiger Reserve. Since land is cheaper near the edges of forests, plantation workers buy these land for building houses. High human population densities in these tea and coffee plantations have led to increasing negative interactions with leopards involving livestock depredation and human injury (Sidhu *et al.*, 2017).



# Kerala



**Figure IV.5:**

*Spatially explicit leopard density (individuals /100 km<sup>2</sup>) modelled from camera traps-based capture-mark-recapture and covariates; Kerala, 2022.*

Kerala is situated at the south-western part of India with a total forest cover of 16,267 km<sup>2</sup> (Forest Survey of India 2017). A total of 10 sites were sampled using camera traps in Kerala that yielded 3,709 photo-captures of 270 individual leopards (Fig. IV.5). The leopard population of the state was estimated at 570 (SE 76), depicting a decline in the leopard population as compared to 650 (SE 28) in 2018 (Jhala *et al.*, 2021). Approximately 63% of the leopard population of Kerala occur outside the Protected Areas of the state.

From 2018 to 2022, Periyar Tiger Reserves showed an increase in leopard population, while the Wayanad and Malayattoor regions experienced a significant decline in leopard population (Table IV.1). Further, Eravikulam National Park, Konni, Ranni and Vazhachal divisions exhibited consistently low leopard densities.

In Kerala, there is a surge in number of human-wildlife conflict cases from 6,022 in 2015-16 to 10,036 in 2021-22 (Issac *et al.*, 2022). There is a rise in incidents of cattle loss and human injury/death caused by leopards (Issac *et al.*, 2022; Karthik, 2020). From 2013-14 to 2018-19, there were a total 547 reported incidents of human-leopard conflict. Total 173 incidents of livestock deaths or injuries (93 cattle, 2 buffalo, 78 goats) were caused by leopards (Kerala State Planning Board, 2022). Most conflict-prone areas were identified as Wayanad, Palakkad, Kannur, Calicut, Thrissur and Malappuram (Jhala *et al.*, 2021). From 2019-20 to 2021-22, most number of human-wildlife conflict incidents were from Wayanad North division, followed by Kannur division and Wayanad South division and the least number of incidents were from the Parambikulam division, followed by Periyar East division and Munnar Wildlife division (Issac *et al.*, 2022).

**Table IV.1: Sampling details and leopard density parameter estimates using spatially explicit capture mark recapture analysis in a likelihood framework for sites in Western Ghats 2022.**

State	Site	Model space (Km <sup>2</sup> )	Camera points	M <sub>t-1</sub>	Best fit model	$\hat{D}$ secr/100 km <sup>2</sup> (SE)	g <sup>0</sup> Male (SE)	g <sup>0</sup> Female (SE)	$\sigma$ Male (SE) (km)	$\sigma$ Female (SE) (km)	Pmix (M:F) (SE)
Goa	Bondla	NA	5	1	NA	NA	NA	NA	NA	NA	NA
Goa	Cotigao	NA	7	2	NA	NA	NA	NA	NA	NA	NA
Goa	Mhadei	418	32	14	$\sigma \sim (.), Pmix \sim (sex), g0 \sim (.)$	7.3 (2.31)	0.055 (0.016)		1.49 (0.21)		0.58:0.42 (0.18)
Goa	Mollem	NA	28	8	NA	NA	NA	NA	NA	NA	NA
Goa	Netravalli	NA	23	5	NA	NA	NA	NA	NA	NA	NA
Karnataka	Ballari	1581	46	30	$\sigma \sim (.), Pmix \sim (sex), g0 \sim (.)$	3.11 (0.64)	0.09 (0.02)		2.67 (0.28)		0.54:0.46 (0.1)
Karnataka	Bandipur TR	1928	612	131	$\sigma \sim (sex), Pmix \sim (sex), g0 \sim (sex)$	9.59 (0.85)	0.03 (0.001)	0.02 (0.001)	1.84 (0.05)	1.27 (0.07)	0.49:0.51 (0.05)
Karnataka	Bengaluru Rural	NA	48	8	NA	NA	NA	NA	NA	NA	NA
Karnataka	Bannerghatta	513	83	30	$\sigma \sim (.), Pmix \sim (sex), g0 \sim (.)$	8.41 (1.56)	0.074 (0.008)		1.75 (0.09)		0.60:0.40 (0.10)
Karnataka	Belagavi	1860	107	41	$\sigma \sim (.), Pmix \sim (sex), g0 \sim (.)$	6.41 (1.44)	0.013 (0.004)		2.17 (0.27)		0.60:0.40 (0.12)
Karnataka	Bhadhravati	1786	120	46	$\sigma \sim (sex), Pmix \sim (sex), g0 \sim (.)$	5.0 (0.78)	0.05 (0.01)		2.92 (0.21)	1.61 (0.13)	0.35:0.65 (0.08)
Karnataka	Bhadra TR	1577	332	112	$\sigma \sim (sex), Pmix \sim (sex), g0 \sim (sex)$	11.31 (1.08)	0.036 (0.003)	0.015 (0.001)	2.03 (0.06)	1.77 (0.06)	0.31:0.69 (0.04)
Karnataka	BRT TR	1076	260	58	$\sigma \sim (sex), Pmix \sim (sex), g0 \sim (sex)$	8.47 (1.14)	0.02 (0.002)	0.011 (0.002)	2.34 (0.12)	1.6 (0.11)	0.29:0.71 (0.06)
Karnataka	Cauvery WLS	2344	448	65	$\sigma \sim (sex), Pmix \sim (sex), g0 \sim (sex)$	5.01 (0.68)	0.02 (0.002)	0.01 (0.002)	3.67 (0.19)	1.84 (0.15)	0.23:0.77 (0.05)
Karnataka	Chikmagalur	793	44	12	$\sigma \sim (.), Pmix \sim (sex), g0 \sim (.)$	4.45 (1.98)	0.04 (0.02)		1.39 (0.35)		0.67:0.33 (0.27)
Karnataka	Davanagere	NA	19	1	NA	NA	NA	NA	NA	NA	NA
Karnataka	Dharwad	NA	26	9	NA	NA	NA	NA	NA	NA	NA
Karnataka	Haliyal	2487	104	27	$\sigma \sim (sex), Pmix \sim (sex), g0 \sim (.)$	4.08 (1.26)	0.03 (0.01)		2.97 (0.37)	1.14 (0.21)	0.16:0.84 (0.07)
Karnataka	Hassan	NA	11	1	NA	NA	NA	NA	NA	NA	NA
Karnataka	Haveri	NA	18	4	NA	NA	NA	NA	NA	NA	NA
Karnataka	Honnavar	1839	125	25	$\sigma \sim (.), Pmix \sim (sex), g0 \sim (.)$	2.77 (0.64)	0.035 (0.009)		1.67 (0.19)		0.73:0.27 (0.11)
Karnataka	Hunsur	NA	36	12	NA	NA	NA	NA	NA	NA	NA

State	Site	Model space (Km <sup>2</sup> )	Camera points	M <sub>t+1</sub>	Best fit model	$\hat{D}$ secr/ 100 km <sup>2</sup> (SE)	g <sup>0</sup> Male (SE)	g <sup>0</sup> Female (SE)	$\sigma$ Male (SE) (km)	$\sigma$ Female (SE) (km)	Pmix (M:F) (SE)
Karnataka	Kalburgi	NA	23	0	NA	NA	NA	NA	NA	NA	NA
Karnataka	Kali TR	3627	399	106	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	6.79 (0.71)	0.022 (0.002)		3.11 (0.13)	1.38 (0.07)	0.22:0.78 (0.04)
Karnataka	Karwar	1594	126	13	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	1.18 (0.35)	0.13 (0.03)			1.28 (0.12)	0.86:0.14 (0.13)
Karnataka	Koppa	NA	49	6	NA	NA	NA	NA	NA	NA	NA
Karnataka	Kudremukh NP	3089	172	58	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	3.8 (0.57)	0.023 (0.004)		2.35 (0.18)		0.58:0.42 (0.09)
Karnataka	Madikeri Territorial	1394	156	14	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	4.4 (1.9)	0.009 (0.005)		1.47 (0.36)		0.37:0.63 (0.17)
Karnataka	Madikeri WLD	1766	169	13	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	2.58 (1.07)	0.01 (0.005)		1.71 (0.41)		0.57:0.43 (0.18)
Karnataka	Mangaluru	NA	57	4	NA	NA	NA	NA	NA	NA	NA
Karnataka	MM Hills	2340	415	51	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	3.59 (0.51)	0.014 (0.002)		2.58 (0.12)		0.90:0.10 (0.06)
Karnataka	Mysore Territorial	NA	35	13	NA	NA	NA	NA	NA	NA	NA
Karnataka	Mysore WLS	174	30	12	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	9.8 (2.98)	0.034 (0.009)		1.98 (0.34)		0.64:0.36 (0.14)
Karnataka	Nagarhole TR	1580	489	97	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	8.85 (0.91)	0.03 (0.002)		1.83 (0.06)	0.96 (0.04)	0.35:0.65 (0.05)
Karnataka	Ramanagara	NA	26	4	NA	NA	NA	NA	NA	NA	NA
Karnataka	Sagara	NA	51	7	NA	NA	NA	NA	NA	NA	NA
Karnataka	Shimoga Territorial	NA	38	4	NA	NA	NA	NA	NA	NA	NA
Karnataka	Shimoga WLD	2647	162	24	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	2.79 (0.66)	0.02 (0.004)		3.34 (0.46)	1.63 (0.23)	0.25:0.75 (0.11)
Karnataka	Sirsi	NA	101	4	NA	NA	NA	NA	NA	NA	NA
Karnataka	Tumkur	NA	24	13	NA	NA	NA	NA	NA	NA	NA
Karnataka	Virajpete	NA	100	2	NA	NA	NA	NA	NA	NA	NA
Karnataka	Yellapura	2722	95	17	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	1.27 (0.39)	0.04 (0.01)		2.35 (0.41)		0.86:0.14 (0.13)
Kerala	Eravikulam NP	NA	28	1	NA	NA	NA	NA	NA	NA	NA
Kerala	Konni	NA	68	10	NA	NA	NA	NA	NA	NA	NA
Kerala	Malayatoor	NA	100	8	NA	NA	NA	NA	NA	NA	NA
Kerala	Parambikulam TR	NA	242	127	$\sigma \sim (\text{sex}), \text{Pmix} \sim (\text{sex}), g_0 \sim (\text{sex})$	13.33 (1.2)	0.07 (0.004)		1.76 (0.04)	1.39 (0.05)	0.33:0.67 (0.04)
Kerala	Periyar TR	2040	396	62	$\sigma \sim (\cdot), \text{Pmix} \sim (\text{sex}), g_0 \sim (\cdot)$	7.19 (1.1)	0.01 (0.001)		2.05 (0.17)		0.85:0.15 (0.07)


State	Site	Model space (Km <sup>2</sup> )	Camera points	M <sub>t+1</sub>	Best fit model	D̂ secr/ 100 km <sup>2</sup> (SE)	g0 Male (SE)	g0 Female (SE)	σ Male (SE) (km)	σ Female (SE) (km)	Pmix (M:F) (SE)
Kerala	Ranni	NA	82	5	NA	NA	NA	NA	NA	NA	NA
Kerala	Silent Valley Landscape	1291	86	12	$\sigma \sim (\cdot), Pmix \sim (\text{sex}), g0 \sim (\cdot)$	2.56 (0.96)	0.011 (0.005)	2.15 (0.43)	NA	NA	0.50:0.50 (0.25)
Kerala	Thattakad	NA	11	0	NA	NA	NA	NA	NA	NA	NA
Kerala	Vazhachal	NA	19	2	NA	NA	NA	NA	NA	NA	NA
Kerala	Wayanad Landscape	2280	281	43	$\sigma \sim (\cdot), Pmix \sim (\text{sex}), g0 \sim (\cdot)$	4.02 (0.71)	0.003 (0.001)	2.5 (0.13)	NA	NA	0.50:0.50 (0.09)
Tamil Nadu	Anamalai TR	2401	496	95	$\sigma \sim (\text{sex}), Pmix \sim (\text{sex}), g0 \sim (\text{sex})$	15.32 (2.99)	0.014 (0.002)	0.004 (0.001)	2.01 (0.13)	1.21 (0.16)	0.12:0.88 (0.03)
Tamil Nadu	Coimbatore	1172	111	32	$\sigma \sim (\cdot), Pmix \sim (\text{sex}), g0 \sim (\cdot)$	4.7 (0.94)	0.035 (0.008)	1.73 (0.16)	NA	NA	0.69:0.29 (0.10)
Tamil Nadu	Erode	1537	368	63	$\sigma \sim (\text{sex}), Pmix \sim (\text{sex}), g0 \sim (\text{sex})$	6.52 (0.84)	0.02 (0.002)	0.01 (0.002)	2.25 (0.11)	1.64 (0.12)	0.38:0.62 (0.07)
Tamil Nadu	Gudalur	720	80	39	$\sigma \sim (\cdot), Pmix \sim (\text{sex}), g0 \sim (\cdot)$	11.9 (2.32)	0.023 (0.005)	1.22 (0.13)	NA	NA	0.57:0.43 (0.11)
Tamil Nadu	Kalakad Mundanthurai TR	1558	228	48	$\sigma \sim (\text{sex}), Pmix \sim (\text{sex}), g0 \sim (\cdot)$	5.61 (0.87)	0.012 (0.002)	2.43 (0.17)	NA	1.46 (0.15)	0.37:0.63 (0.08)
Tamil Nadu	Kanyakumari	NA	105	5	NA	NA	NA	NA	NA	NA	NA
Tamil Nadu	Kodaikanal	560	97	12	$\sigma \sim (\cdot), Pmix \sim (\text{sex}), g0 \sim (\cdot)$	5.13 (1.63)	0.012 (0.004)	1.53 (0.22)	NA	NA	0.33:0.67 (0.19)
Tamil Nadu	Mudumalai TR	1903	394	120	$\sigma \sim (\text{sex}), Pmix \sim (\text{sex}), g0 \sim (\text{sex})$	11.56 (1.08)	0.027 (0.002)	0.018 (0.003)	2.11 (0.07)	1.04 (0.06)	0.30:0.70 (0.04)
Tamil Nadu	Mukurthy NP	685	50	9	NA	NA	NA	NA	NA	NA	NA
Tamil Nadu	Nellai	975	80	19	$\sigma \sim (\cdot), Pmix \sim (\text{sex}), g0 \sim (\text{sex})$	3.35 (0.81)	0.008 (0.002)	0.039 (0.014)	3.12 (0.27)	NA	0.76:0.24 (0.09)
Tamil Nadu	Nilgiri	2133	328	121	$\sigma \sim (\text{sex}), Pmix \sim (\text{sex}), g0 \sim (\text{sex})$	13.14 (1.39)	0.201 (0.43)	0.028 (0.005)	1.88 (0.1)	0.79 (0.06)	0.30:0.70 (0.05)
Tamil Nadu	Sathyamangalam TR	2700	704	103	$\sigma \sim (\text{sex}), Pmix \sim (\text{sex}), g0 \sim (\cdot)$	7.05 (0.75)	0.011 (0.001)	2.62 (0.11)	NA	1.2 (0.07)	0.30:0.70 (0.04)
Tamil Nadu	Srivilliputhur	1219	264	44	$\sigma \sim (\text{sex}), Pmix \sim (\text{sex}), g0 \sim (\cdot)$	6.42 (1.03)	0.01 (0.001)	2.35 (0.19)	NA	1.74 (0.18)	0.45:0.55 (0.11)
Tamil Nadu	Meghamalai	1490	344	39	$\sigma \sim (\cdot), Pmix \sim (\text{sex}), g0 \sim (\cdot)$	4.63 (0.77)	0.01 (0.001)	1.97 (0.14)	NA	NA	0.5:0.5 (0.11)

**SE:** Standard error, **D̂SECR:** Density estimate from Maximum Likelihood based spatially explicit capture recapture, **σ** : Spatial scale of detection function, **g0:** Magnitude (intercept) of detection function, **Pmix:** Detection corrected estimate of proportion of female and male

# Conservation Implications

The leopard is the most widely distributed large carnivore in the Western Ghats landscape, and involved in the most of the large carnivore-human conflict cases. The leopard populations in the Western Ghats landscape are facing the wrath of rapid development, habitat loss, and retaliatory killing by humans. Although the latest figures of the

leopard population estimates depicted slight increase in the landscape, local declines were observed in some of the strongholds (e.g., Wayanad). It is crucial to investigate the population status and collate further information on poaching, human-leopard conflict, and status of co-predators to understand population dynamics of leopards and make appropriate and timely conservation interventions. Studies focused on land-tenure systems of leopards (involving radio-telemetry) and demographic parameters of leopards from different parts of the landscape will provide valuable insights into the lesser known aspects of species ecology, thus help making knowledge-based conservation actions.





©Vinay Venugopal





# V: North East Hills and Brahmaputra Flood Plains

Deb Ranjan Laha, Shikha Bisht, Ujjwal Kumar, Vishnupriya Kolipakam, Swati Saini, Richard S. Sangma, Vaishnavi Gusain, Dhruv Jain, Omkar Nar, Anshuman Gogoi, Monika Saraswat, Rajendra Garawad, Satya P. Yadav, Yadvendradev V. Jhala and Qamar Qureshi



©Tanjil Tamuli

## North East Hills and Brahmaputra Flood Plains

Eastern Himalaya and its foothills including the north Bengal *Dooars*, and fertile floodplains of Brahmaputra and Barak along with north east Hills act as Indo-Malayan biodiversity hotspot and home to several endangered and endemic faunas. However, due to geo-political and developmental reasons, this landscape undergone several land-use changes which led to sever loss of natural habitats across the landscape (Jhala *et al.*, 2020).


Leopards are present across the entire landscape, however, in this report estimate of leopard is limited to the sampled forested area only. Spread of phase I sampling was sparse in Arunachal Pradesh, Assam and Mizoram and restricted mostly in the Tiger Reserve. The entire landscape is mosaic of forested habitat, agriculture and plantations, and several other human land use features thus provide excellent cover for leopards which are quite adaptable with regards to their habitat preference and dietary need (Jhala *et al.*, 2021) and often found in areas bordering human dominated hab-


itats. This causes significant human leopard conflict in this landscape (Marker and Sivamani 2009, Bhattacharjee and Parthasarathy 2013, Kshetry *et al.*, 2017, Naha *et al.*, 2018). Despite being present all across the landscape, leopards are least studied species in the landscape and there are very few studies available on population estimates of leopards since last decade (Jhala *et al.*, 2021). Jhala *et al.*, 2021 published the baseline population estimates of leopard at landscape and individual site level, however, this too was a snapshot of the population of leopard in this landscape.

## Leopard Occupancy, Population Extent and Abundance

Leopard signs are recorded from north Bengal *Dooars*, Assam, Arunachal Pradesh, Mizoram and Nagaland. However, except north Bengal *Dooars* where Phase I sampling was carried out outside Protected Areas as well, leopard signs obtained are very sporadic in other states owing to the poor Phase I sampling. In Assam, especially in upper Assam and areas like Karbi-Anglong and western Assam where

**Figure V.1:** Change in leopard distribution in North East Hills and Brahmaputra Flood Plains landscape from 2018 to 2022.





**Figure V.2:** Leopard distribution and density for North East Hills and Brahmaputra Flood Plains landscape, 2022.

leopards were detected in 2018 had not been sampled in 2022 (Fig. V.1). There is an increase in sampling effort in north Bengal *Dooars*, Arunachal Pradesh and Nagaland.

A total of 250 cells (100 km<sup>2</sup>) were sampled in 2022 in this landscape, out of which leopard presence were recorded from 53%(133) cells. Almost one third of the total cells in 2022 were not sampled in 2018 and have recorded leopard presence in 2022 (Fig. V.1).

A total of 15 sites in this landscape were sampled using camera trap based mark-recapture method, and leopard images were obtained from 11 sites. Due to inadequate spatial and temporal photo-captures, density estimates could not be assessed for Namdapha Tiger Reserve of Arunachal Pradesh, Neora valley National Park and Mahananda Wildlife Sanctuary of West Bengal. Phase I sampling in Arunachal Pradesh was restricted mainly to Tiger Reserves, thus leopard


density could not be extrapolated outside Tiger Reserves in the states (Table V.1). In addition, we could not estimate leopard population for Nagaland and Mizoram as no leopard positive scat and/ or leopard images were obtained from these two states.

The leopard population of this landscape was estimated to be 349 (SE 42) as compared to 141 (SE 26) in 2018. 228 individual leopards were identified from 4614 photo-captures. The increase in the population estimate is mainly because of sampling artefact, as Phase I and III sampling effort increased significantly in north Bengal *Dooars* in 2022. The figure V.2 depicts leopard density in the landscape, however, in Nagaland, Mizoram and parts of Arunachal Pradesh and Assam the low density represents only the presence of leopard signs in sampled forested areas.



©Gurinderjit Singh

# Arunachal Pradesh



**Figure V.3:**  
Leopard distribution and density for Arunachal Pradesh, 2022.

Camera trap based mark recapture sampling was done in four sites in the state namely Kamlang, Namdapha and Pakke Tiger Reserves and Dibrang Wildlife Sanctuary. However, leopard images were obtained only from Namdapha and Pakke Tiger Reserves. Leopard presence was not confirmed from Kamlang and Dibrang, either from camera trap or through scat DNA.


News report of a straying leopard being rescued from Wakkro town in Lohit district adjacent to Kamlang Tiger Reserve in 2022 (The Arunachal Times, 2022) suggests the presence of the leopard around Kamlang Tiger Reserve.

A total of 41 individual leopards were identified from 796 photo-captures in Arunachal Pradesh of which four leop-

ards were common between Arunachal Pradesh (Pakke Tiger Reserve) and Assam (Nameri Tiger Reserve). Estimated leopard population of Arunachal Pradesh was 42 (SE 10) which is an underestimation as leopard population was estimated only from the camera-trapped sites. There is a substantial increase in the population density of leopards in Pakke since 2018 (Jhala *et al.*, 2021) (Table V.1).

Moderate to high density of leopards can be observed in Pakke Tiger Reserve. As several developmental projects are coming up in Arunachal Pradesh (Qureshi *et al.*, 2013), there is an urgent need of sampling outside Tiger Reserves in the state to understand the spatial extent and abundance of leopards and other wildlife.

# Assam



**Figure V.4:**


*Leopard distribution and density for Assam, 2022.*

Five sites were camera trapped in using mark-recapture framework in the state (Table V.1) and only one site (Nagaon Wildlife Division) had no leopard photo-capture. A total of 726 photo-captures yielded 62 individual leopards and leopard population was estimated at 74 (SE 11). Four leopards of Nameri Tiger Reserve were common with neighbouring Pakke Tiger Reserve in Arunachal Pradesh. An individual leopard was photo-captured from Orang Tiger Reserve for the very first time (Table V.1). Population estimate of leopards is certainly an underestimation, as being an adaptable feline, leopards are reported from almost every corner of the state. In addition, studies have reported presence of leopard from Raimona National Park and Ripu Reserve Forest of western Assam (Nath *et al.*, 2021, Nath *et al.*, 2023) where Phase I and III sampling was not carried out. The entire Manas- Buxa-Royal Manas complex acts as a single population block thus maintaining habitat integrity in this complex is crucial. Low to moderate density of leopards can be observed in sampled forests of Assam, with few parts of Manas and Kaziranga having high density. Population estimate of leopards is limited to the Tiger Reserves, and leopard density could not be extrapolated in the state owing to the extremely poor sampling spread of Phase I. Many of the forest divisions in upper As-

sam, western Assam and areas of Karbi Anglong where leopard presence was recorded earlier in 2018, were not sampled in 2022. Although, leopards are recorded from most parts of the state, only a few studies have been carried out to assess the status and spatial extent of leopard (Bora *et al.*, 2013, Harihar *et al.*, 2020, Jhala *et al.*, 2021). Density of leopard in Kaziranga, Manas and Nameri remain stable in comparison to 2018 (Jhala *et al.*, 2021) (Table V.1).

Due to the habitat being a mosaic of forests, tea plantations and agricultural lands which serve as excellent cover for leopards, growing human leopard conflict is a major concern towards conservation of leopard in the state. In addition, increased traffic in several state and national highways also causes mortality of leopards in the state. However, a thorough Phase I sampling in the forested habitat of the state along with camera trapping exercise in Protected Areas are essential to identify the spatial extent and status of the leopard in the state. This would also help wildlife managers and policy makers to plan and mitigate negative human leopard interaction in the state.

# West Bengal (North Bengal *Dooars*)



**Figure V.5:**

*Leopard distribution and density for West Bengal (North Bengal Dooars), 2022.*

North Bengal *Dooars* is an extension of Indo-Gangetic alluvial floodplains and situated along the foothills of eastern Himalayas. For the very first time, an extensive area including forested habitats outside Protected Areas in this landscape have been systematically sampled to collect Phase I data. In addition, a total of five sites were sampled in camera trap based mark-recapture framework and leopard photo-captures were recorded in all the sites (Table V.1). A total of 3,092 photo-captures of leopard yielded 129 individuals. However, due to low detection and inadequate spatial and temporal photo-captures of individual leopards in Neora valley National Park and Mahananda Wildlife Sanctuary, Spatially Explicit Capture Recapture (SECR) based density estimates was not possible.

Despite leopard being present across north Bengal *Dooars*, there are only two studies (Borthakur *et al.*, 2021, Jhala *et al.*, 2021) available, which have reported the status of leopard population in north Bengal *Dooars*. As per Borthakur *et al.*, 2021, population of leopard was estimated at 111 obtained from scat based DNA sampling in north Bengal *Dooars*, whereas Jhala *et al.*, 2021 provided a snapshot of baseline data from mark-recapture based camera trap data obtained from three Protected Areas namely Buxa Tiger Reserve, Gorumara and Jaldapara Nation-

al Parks. Leopard population of north Bengal *Dooars* estimated at 233 (SE 21) which is more than two fold as compared to 2018 (Jhala *et al.*, 2021) (Table I.3). However, this could be an artefact of sampling, as the sampling effort has increased and Phase I data was collected outside the camera trap sites as well, which was not the case in earlier cycles. Leopard signs have been obtained from areas which were not sampled in 2018. Density estimates of leopard remained stable for Jaldapara National Park (Table V.1), however, there was a substantial increase in leopard population and density estimates of Buxa Tiger Reserve and Gorumara National Park as compared to 2018 (Jhala *et al.*, 2021) (Fig V.1).

Like Assam, large forested tracks of north Bengal *Dooars* had been exploited during colonial era for cultivation tea and revenue generation by timber extraction. Majority of the human population living in tea estates are marginal workers brought from Chotanagpur plateau region during colonial period and rear livestock for livelihood. This has resulted in fragmentation of natural habitats situated in the mosaic of tea gardens and several other human land use features. Leopards, however, with a flexible and broader niche and dietary preference have adapted in this fragmented landscape as the mosaic habitat characteristics provide ample cover for leopards. With increasing human

## West Bengal (North Bengal *Dooars*)

population, human leopard conflict in the landscape has escalated and acts as major challenge for conservation of leopard in north Bengal *Dooars* (Vyas and Sengupta 2014). A study by Naha *et al.*, 2018 reported that 97% of human injuries are resulted by leopard attack in North Bengal as compared to 60% in Pauri Garhwal district of Uttarakhand. Much of the literature from north Bengal *Dooars* addressed human leopard conflict, dietary preference, and niche preference (Kshetry *et al.*, 2017, Kshetry *et al.*, 2018, Naha *et al.*, 2018, Naha *et al.*, 2020, Naha *et al.*, 2021, Borthakur *et al.*, 2021). Major portion of the diet (~65%) of leopards in north Bengal *Dooars* comprised of livestock which complement wild prey species (Borthakur *et al.*, 2021, Kshetry 2023) and have higher livestock depredation risk (Naha *et al.*, 2020).

North Bengal *Dooars* are connecting link to the peninsular India (Qureshi *et al.*, 2023) and hence numerous linear infrastructures traverse north Bengal *Dooars* to maintain geo-political integrity with neighbouring countries, which further complicates the scenario of leopard conservation in this landscape. Especially, National Highway NH31 and railway line between Siliguri- Alipurduar junctions cuts across many forested areas and at least three Protected Areas (Mukherjee *et al.*, 2019) and leads to accidents with many wild animals and which need appropriate mitigation measures. An active management plan with community perspectives should be prepared to minimize human-leopard conflict in north Bengal *Dooars*.

## Conservation Implications

Although there is an increase in leopard population in this landscape as compared to earlier cycles. Majority of leopard population is recorded from north Bengal *Dooars*, which was due to increased sampling effort. Due to extremely low abundance of tiger (Qureshi *et al.*, 2023), leopard serve as apex predator in the Protected Areas of north Bengal *Dooars*. Leopard density in North East Hills and Brahmaputra Flood Plains landscape varies from 1.94 (SE 0.53) leopards/ 100 km<sup>2</sup> in Kaziranga Tiger Reserve of Assam to 15.12 (SE 2.75) leopards/ 100 km<sup>2</sup> in Gorumara National Park of West Bengal (Table V.1). Also, for the very first time a leopard was photo-captured in Orang Tiger Reserve. Arunachal Pradesh also has recorded increase in leopard population in sampled area (Jhala *et al.*, 2021). Kamlang Tiger Reserve has not recorded any leopard in current cycle of All India Tiger Estimation 2022. Leopard presence was also not confirmed from Dampa Tiger Reserve (Mizoram) from either camera trap image or scat. A thorough sampling throughout these Tiger Reserves is crucial to understand the status and spatial extent of large carnivores. North Bengal *Dooars* is situated amidst mosaic of human land-use patterns and forested patches, managerial efforts are needed to carefully execute the process to minimize human-leopard conflict.

Buxa Tiger Reserve is preparing for ambitious tiger re-introduction program, and with its growing population of leopards this should be taken up with caution. Since Buxa harbours low abundance of wild prey species (Jhala *et al.*, 2020), stocking of wild prey and prey augmentation procedures should take into account other large carnivores of this area.

This report, however, provides an extended baseline information of status of leopards in this landscape. In addition, systematic and thorough sampling for covariates through Phase I survey in the floodplains and foothills of eastern Himalayas and camera trapping and/ or scat collection in the North Eastern hills are crucial to identify conservation priority areas and to mitigate negative human leopard interactions.



**Table V.1:** Sampling details and leopard density parameter estimates using spatially explicit capture mark recapture analysis in a likelihood framework for sites in the North East Hills and Brahmaputra Flood Plains landscape, 2022.

State	Site	Model Space (km <sup>2</sup> )	Camera traps	M <sub>t+1</sub>	Best Fit Model	$\hat{D}_{SECR}/100 \text{ km}^2$ (SE)	g0 Female (SE)	g0 Male (SE)	$\sigma$ Female (SE) (km)	$\sigma$ Male (SE) (km)	Pmix (SE)
Arunachal Pradesh	Namdapha TR	NA	117	5	NA	NA	NA	NA	NA	NA	NA
Arunachal Pradesh	Pakke TR	1040	135	36	$\sigma \sim (\cdot), P_{mix} \sim (\text{sex}), g0 \sim (\cdot)$	4.2 (0.7)	0.03 (0.003)	0.03 (0.003)	2.55 (0.11)	2.55 (0.11)	0.40:0.60 (0.1)
Assam	Nameri TR	579	73	12	$\sigma \sim (\cdot), P_{mix} \sim (\text{sex}), g0 \sim (\cdot)$	2.66 (0.8)	0.03 (0.003)	0.03 (0.003)	2.55 (0.11)	2.55 (0.11)	0.20:0.80 (0.1)
Assam	Orang TR	NA	39	1	NA	NA	NA	NA	NA	NA	NA
Assam	Kaziranga TR	747	230	14	$\sigma \sim (\cdot), P_{mix} \sim (\text{sex}), g0 \sim (\cdot)$	1.94 (0.5)	0.005 (0.001)	0.005 (0.001)	4.53 (0.34)	4.53 (0.34)	0.70: 0.30 (0.1)
Assam	Manas TR	1052	267	35	$\sigma \sim (\text{sex}), P_{mix} \sim (\text{sex}), g0 \sim (\text{sex})$	4.19 (0.7)	0.006 (0.001)	0.02 (0.002)	2.21 (0.19)	2.87 (0.18)	0.70:0.30 (0.1)
West Bengal	Buxa TR	802	162	61	$\sigma \sim (\cdot), P_{mix} \sim (\text{sex}), g0 \sim (\cdot)$	9.17 (1.2)	0.03 (0.002)	0.03 (0.002)	2.01 (0.06)	2.01 (0.06)	0.60:0.40 (0.1)
West Bengal	Jaldapara NP	345	75	30	$\sigma \sim (\cdot), P_{mix} \sim (\text{sex}), g0 \sim (\cdot)$	13.84 (2.8)	0.02 (0.005)	0.02 (0.005)	1.13 (0.12)	1.13 (0.12)	0.50:0.50 (0.1)
West Bengal	Gorumara NP	217	49	31	$\sigma \sim (\cdot), P_{mix} \sim (\text{sex}), g0 \sim (\cdot)$	15.12 (2.8)	0.04 (0.004)	0.04 (0.004)	1.96 (0.1)	1.96 (0.1)	0.40: 0.60 (0.1)
West Bengal	Neora Valley NP	NA	21	1	NA	NA	NA	NA	NA	NA	NA
West Bengal	Mahananda WLS	NA	22	6	NA	NA	NA	NA	NA	NA	NA

SE: Standard error, D<sup>SECR</sup>: Density estimate from Maximum Likelihood based spatially explicit capture recapture, : Spatial scale of detection function, g0: Magnitude (intercept) of detection function, Pmix: Detection corrected estimate of proportion of female and male





# References

- Akaike, H. 2011. Akaike's Information Criterion. In: Lovric, M. (eds) International Encyclopedia of Statistical Science. Springer, Berlin, Heidelberg. [https://doi.org/10.1007/978-3-642-04898-2\\_110](https://doi.org/10.1007/978-3-642-04898-2_110)
- Athreya, V. 2012. Conflict resolution and leopard conservation in a human dominated landscape. Ph.D. thesis, Manipal University, India, 146 p.
- Athreya, V., Odden, M., Linnell, J.D.C. and Karanth, K.U. 2011. Translocation as a tool for mitigating conflict with leopards in human-dominated landscapes of India. *Conservation Biology*, 25, 133–141.
- Athreya, V., Odden, M., Linnell, J.D.C., Krishnaswamy, J. and Karanth, U. 2013. Big cats in our backyards: persistence of large carnivores in a human dominated landscape in India. *PLoS One*, 8, e57872.
- Athreya, V., Odden, M., Linnell, J.D.C., Krishnaswamy, J. and Karanth, K.U. 2014. A cat among the dogs: leopard *Panthera pardus* diet in a human-dominated landscape in western Maharashtra, India. *Oryx*, 50(1), 156-162.
- Athreya, V., Srivathsa, A., Puri, M., Karanth, K.K., Kumar, N.S. and Karanth, K.U. 2015. Spotted in the news: using media reports to examine leopard distribution, depredation, and management practices outside Protected Areas in Southern India. *PLoS One*, 10(11), e0142647.
- Bailey, T.N. 1993. The African leopard: ecology and behavior of a solitary felid. Columbia University Press, New York, 429 p.
- Bargali, H.S. and Ahmed, T. 2018. Patterns of livestock depredation by tiger (*Panthera tigris*) and leopard (*Panthera pardus*) in and around Corbett Tiger Reserve, Uttarakhand, India. *PLoS One*, 13(5), e0195612.
- Bhatt, S., Biswas, S., Karanth, K., Pandav, B. and Mondol, S. 2020. Genetic analyses reveal population structure and recent decline in leopards (*Panthera pardus fusca*) across the Indian subcontinent. *Peer J*, 8, e8482.
- Bhattacharjee, A., & Parthasarathy, N. 2013. Coexisting with large carnivores: a case study from Western Duars, India. *Human dimensions of wildlife*, 18(1), 20-31.
- Bista, Ashish., Chanchani, Pranav., Pandey, Ramesh., Hasan, Dabeer, Gupta, Mudit. and Pathak, Sanjay. 2019. Status of Common Leopard in Katerniaghat Wildlife Division. Uttar Pradesh Forest Department and World Wide Fund for Nature-India.
- Biswas, S., Bhatt, S., Sarkar, D., Talukdar, G., Pandav, B. and Mondol, S. 2020. Assessing tiger corridor functionality with landscape genetics and modelling across Terai-Arc Landscape, India. *bioRxiv*.
- Blanford, W.T. 1891. Fauna of British India. Taylor and Francis, London.
- Borah, J., Sharma, T., Das, D., Rabha, N., Kakati, N., Basumatary, A., Ahmed, M. F. and Vattakaven, J. 2014. Abundance and density estimates for common leopard *Panthera pardus* and clouded leopard *Neofelis nebulosa* in Manas National Park, Assam, India. *Oryx*, 48 (1), 149–155.
- Borchers, D.L. and Efford, M.G. 2008. Spatially explicit maximum likelihood methods for capture recapture studies. *Biometrics*, 64 (2), 377-385.
- Borthakur, U., Das, P.K., Deb, M., Maheen, R., Neog, A., Das, A.K., Baruah, B., Singha Roy, R., Kalita, J., Biswas, A. and Kurmi, R.P. 2017. Leopard (*Panthera pardus*) ecology in North Bengal including population estimation, population dynamics, distribution, habitat use pattern and human leopard conflicts. Technical Report, Aaranyak, Guwahati, India.
- Borthakur, U., Das, P.K., Deb, M., Maheen, R., Neog, A., Das, A.K., Barua, B. and Biswas, A. 2021. Leopard (*Panthera pardus*) ecology in North Bengal including population estimation, distribution, habitat use pattern and human-leopard conflicts. Aaranyak, Technical Report, Pg. 1-87.
- Caro, T.M. 2003. Umbrella species: critique and lessons from East Africa. *Animal Conservation*, 6, 171-181.
- Champion, F.W. 1927. With a camera in tiger-land. Chatto and Windus, London, xviii+228 p.
- Champion, H. and Seth, S.K. 1968. A revised survey of the forest types of India. Daya Publishing House, Delhi, 404 p.
- Chanchani, P., Bista, A., Warriar, R., Nair, S., Sharma, R., Hassan, D., & Gupta, M. 2014. Status and conservation of tigers and their prey in the Uttar Pradesh Terai. WWF-India, New Delhi.
- Chapron, G., Miquelle, D.G., Lambert, A., Goodrich, J.M., Legendre, S. and Clobert, J. 2008. The impact on tigers of poaching versus prey depletion. *Journal of Applied Ecology*, 45, 1667–1674.
- Chatterjee, S. 2008. Biodiversity conservation issues of Northeast India. *The International Forestry Review*, 10(2), 315-324.
- Chaudhary, R., Zehra, N., Musavi, A. and Khan, J.A. 2020. Spatio-temporal partitioning and coexistence between leopard (*Panthera pardus fusca*) and Asiatic lion (*Panthera leo persica*) in Gir protected area, Gujarat, India. *PLoS One*, 15(3), e0229045.
- Chauhan, D.S. and Goyal, S.P. 2001. A study on distribution, relative abundance and food habits of Leopard (*Panthera pardus*) in Garhwal Himalayas. Technical Report (December 1999 – July 2000), Wildlife Institute of India, Dehradun.
- Chauhan, D.S., Harihar, A., Goyal, S.P., Qureshi, Q., Lal, P. and Mathur, V.B. 2005. Estimating leopard population using camera traps in Sariska Tiger Reserve. Wildlife Institute of India, Dehradun.
- Chauhan, K., Srivathsa, A., & Athreya, V. 2021. Assessing spatio-temporal patterns of human-leopard interactions based on media reports in northwestern India. *Journal of Threatened Taxa*, 13(6), 18453-18478.
- Cheema, G.S., Anand, S., Qureshi, Q. and Jhala, Y.V. 2018. CaTRAT: Camera Trap Data Repository and Analysis Tool. Wildlife Institute of India, Dehradun and Indraprastha Institute of Information Technology, New Delhi. Available at [https://wii.gov.in/tiger\\_monitoring\\_software](https://wii.gov.in/tiger_monitoring_software).
- Chellam, R. 1993. Ecology of the Asiatic lion (*Panthera leo persica*). Ph.D. Thesis, Saurashtra University, Rajkot, India, 170 p.
- Crall, J. P., Stewart, C., Berger-Wolf, T., Rubenstein, D., & Sundaresan, S. 2013. HotSpotter — Patterned species instance recog-

- nition. In Proceedings of IEEE Workshop on Applications of Computer Vision. <https://doi.org/10.1109/WACV.2013.6475023>
- Daniel, J.C. 1996. The leopard in India- a natural history. Natraj Publishers. Dehradun, 228 p.
  - Datta A., Anand M.O. and Naniwadekar R. 2008. Empty forests: Large carnivore and prey abundance in Namdapha National Park, Northeast India. *Biological Conservation*, 141, 1429-1435.
  - Deivanayaki, M., & Ezhilarasi, N. 2019. Human–Carnivore conflict in Anamalai Tiger Reserve, Tamil Nadu, Southern Western Ghats. *J. Sci*, 12(3), 113-119.
  - Dhanwatey, H., Crawford, J., Abade, L., Dhanwatey, P., Nielsen, C., and Sillero-Zubiri, C. 2013. Large carnivore attacks on humans in central India: A case study from the Tadoba-Andhari Tiger Reserve. *Oryx*, 47(2), 221-227.
  - Dutta, T., Sharma, S., Maldonado, J.E., Wood, T.C. and Seidensticker, J. 2012. A reliable method for individual identification and gender determination of wild leopards (*Panthera pardus fusca*) using non-invasive samples. *Conservation Genetics Resources*, 4, 665–667.
  - Dutta, T., Sharma, S., Maldonado, J.E., Wood, T.C., Panwar, H.S. and Seidensticker, J. 2013. Fine-scale population genetic structure in a wide-ranging carnivore, the leopard (*Panthera pardus fusca*) in central India. *Diversity and Distributions*, 19(7), 760-771.
  - Dutta, T., Sharma, S., Maldonado, J.E., Wood, T.C., Panwar, H.S. and Seidensticker, J. 2013. Gene flow and demographic history of leopards (*Panthera pardus*) in the central Indian highlands. *Evolutionary Applications*, 6, 949–959.
  - Edgaonkar, A. 2008. Ecology of the leopard (*Panthera pardus*) in Bori Wildlife Sanctuary and Satpura National Park, India. Ph.D. thesis, University of Florida, USA. 135 p.
  - Efford, M.G. 2011. Estimation of population density by spatially explicit capture–recapture analysis of data from area searches. *Ecology*, 92, 2202–2207.
  - Efford, M.G. 2015. secr: spatially explicit capture-recapture models. - R package version 2.9.4. <http://CRAN.R-project.org/package=secur>.
  - Elvidge, C.D., Baugh, K., Zhizhin, M., Hsu, F.C. and Ghosh, T. 2017. VIIRS night-time lights. *International Journal of Remote Sensing*, 38 (21), 5860-5879.
  - Fick, S.E. and Hijmans, R.J. 2017. WorldClim 2: new 1km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, 37 (12), 4302-4315.
  - Finn, F. 1929. Mammalia of India. Thaker, Spink and Co., Bombay, 347 p.
  - Forest Survey of India, 2017. India State of forest-2017. Ministry of Environment, Forest and Climate Change
  - Forest Survey of India, 2021. India State of forest-2021. Ministry of Environment, Forest and Climate Change
  - Gubbi, S., Kolekar, A., & Kumara, V. 2020. Policy to on-ground action: Evaluating a conflict policy guideline for leopards in India. *Journal of International Wildlife Law & Policy*, 23(2), 127-140.
  - Gubbi, S., Nagashettihalli, H., Bhat, R., Poornesha, H.C., Anoop, A. and Madhusudan, M.D. 2017. Ecology and conservation of leopards in protected and multiple use forests in Karnataka. Nature Conservation Foundation, Mysore, India.
  - Gubbi, S., Nagashettihalli, H., Suthar, S. and Menon, A.M. 2019. Report on monitoring of leopards at Biligiri Rangaswamy Temple Tiger Reserve in Karnataka. Nature Conservation Foundation, Mysore, India.
  - Gubbi, S., Poornesha, H.C., Daithota, A. and Nagashettihalli, H. 2014. Roads emerging as a critical threat to leopards in India. *Cat News*, 60, 30–31.
  - Gubbi, S., Ramesh, S., Menon, A.M., Girish, M.N. and Poornesha, H.C. 2020. The lone wolf: new distribution update of the Indian grey wolf (*Canis lupus pallipes*) in southern India. *Canid Biology and Conservation*, 22(6), 21-24.
  - Gubbi, S., Sharma, K. and Kumara, V. 2020. Every hill has its leopard: patterns of space use by leopards (*Panthera pardus*) in a mixed use landscape in India. *Peer J*, 8, e10072.
  - Harihar, A., Lahkar, D., Singh, A., Das, S. K., Ahmed, M. F., & Begum, R. H. 2021. Population density modelling of mixed polymorphic phenotypes: an application of spatial mark-resight models. *Animal Conservation*, 24(4), 709-716.
  - Harihar, A., Pandav, B. and Goyal, S.P. 2009. Density of leopards (*Panthera pardus*) in the Chilla Range of Rajaji National Park, Uttarakhand, India. *Mammalia*, 73, 68–71.
  - Harihar, A., Pandav, B. and Goyal, S.P. 2011. Responses of leopard *Panthera pardus* to the recovery of a tiger *Panthera tigris* population. *Journal of Applied Ecology*, 48(3), 806-814.
  - Hayward, M., Henschel, P., O'Brien, J., Hofmeyr, M., Balme, G. and Kerley, G. 2006. Prey preferences of the leopard (*Panthera pardus*). *Journal of Zoology*, 270, 298-313.
  - Henle, K., Lindenmayer, D.B., Margules, C.R., Saunders, D.A. and Wissel, C. 2004. Species survival in fragmented landscapes: where are we now? *Biodiversity and Conservation*, 13(1), 1-8.
  - Herald Goa 2024. Man-animal conflict surfaces again. <https://www.heraldgoa.in/Edit/Mananimal-conflict-surfaces-again/189814>
  - Herald Goa 2024 No conflict: Simple case of insensitive humans destroying animal habitats. <https://www.heraldgoa.in/Goa/No-conflict-Simple-case-of-insensitive-humans-destroying-animal-habitats/216182>
  - Hiby, L., Lovell, P., Patil, N., Kumar, N.S., Gopalaswamy, A.G. and Karanth, K.U. 2009. A tiger cannot change its stripes: using a three-dimensional model to match images of living tigers and tiger skins. *Biology Letters*, 5 (3), 383 - 386.
  - Hoffmann, M. and Atickem, A. 2019. *Canis lupaster*. The IUCN Red List of Threatened Species 2019, e.T118264888A118265889.
  - Hoffmann, M., Arnold, J., Duckworth, J.W., Jhala, Y., Kamler, J.F. and Krofel, M. 2018. *Canis aureus*. The IUCN Red List of Threatened Species 2018, e.T118264161A46194820.
  - India Today (2021) Bengal: Car runs over leopard in Alipurduar, driver tries to flee with carcass <https://www.indiatoday.in/india/story/bengal-car-runs-over-leopard-in-alipurduar-driver-tries-to-flee-with-carcass-1808220-2021-05-28>

- Indian Express (2023) Goa: Leopard trapped in snare dies, probe on. <https://indianexpress.com/article/india/goa-leopard-trapped-in-snare-dies-probe-on-8693344/>
- Issac, C. J., & Prasad, G. (2022). Compensation Records as an Indirect Method of Assessing Human-wildlife Conflict: A Case Study From Kerala, India.
- Jacobson AP, Gerngross P, Lemeris Jr. JR, Schoonover RF, Anco C, Breitenmoser-Würsten C, Durant SM, Farhadinia MS, Henschel P, Kamler JF, Laguardia A, Rostro-García S, Stein AB, Dollar L. 2016. Leopard (*Panthera pardus*) status, distribution, and the research efforts across its range. *PeerJ* 4:e1974 <https://doi.org/10.7717/peerj.1974>
- Jhala, H.Y., Qureshi, Q., Jhala, Y.V. and Black, S.A., 2021. Feasibility of reintroducing grassland megaherbivores, the greater one-horned rhinoceros, and swamp buffalo within their historic global range. *Scientific reports*, 11(1), pp.1-15.
- Jhala, Y.V., Qureshi, Q., Yadav, S.P. 2021. Status of leopards, co-predators, and megaherbivores in India, 2018. National Tiger Conservation Authority, government of India, New Delhi and Wildlife Institute of India, Dehradun, ISBN-81-85496-56-0.
- Jhala, Y.V., Qureshi, Q. and Gopal, R. (Eds.) 2015. The status of tigers in India, 2014. National Tiger Conservation Authority, New Delhi and Wildlife Institute of India, Dehradun, 456 p.
- Jhala, Y.V., Qureshi, Q. and Gopal, R. 2011. Can the abundance of tigers be assessed from their signs? *Journal of Applied Ecology*, 48, 14-24.
- Jhala, Y.V., Qureshi, Q. and Gopal, R. 2017. Field Guide: Monitoring tigers, co-predators, prey and their habitats. Technical Publication of National Tiger Conservation Authority, New Delhi and the Wildlife Institute of India, Dehradun. Technical Report TR-2017/012.
- Jhala, Y.V., Qureshi, Q. and Nayak, A.K. (eds) 2020. Status of tigers, copredators and prey in India, 2018. National Tiger Conservation Authority, Government of India, New Delhi, and Wildlife Institute of India, Dehradun ISBN No. 81-85496-50-1
- Jhala, Y.V., Qureshi, Q. and Yadav, S.P., 2020. Status of leopards in India, 2018. National Tiger Conservation Authority, Government of India, New Delhi, and Wildlife Institute of India. Dehradun. Technical Report TR/2020/16.
- Jhala, Y.V., Qureshi, Q., Nayak, A.K., 2019. Status of tigers, co-predators and prey in India 2018 (Summary No. ISBN 81-85496-50-1). National Tiger Conservation Authority, Government of India, New Delhi, and Wildlife Institute of India Dehradun, Dehradun.
- Johnsingh, A.J.T. 1983. Large mammalian prey-predators in Bandipur. *Journal of the Bombay Natural History Society*, 80, 1-57.
- Johnsingh, A.J.T., 1992. Prey selection in three large sympatric carnivores in Bandipur.
- Johnsingh, A.J.T., Ramesh, K., Qureshi, Q., David, A., Goyal, S.P., Rawat, G.S., Rajapandian, K. and Prasad, S. 2004. Conservation status of tiger and associated species in the Terai Arc Landscape, India. Wildlife Institute of India, Dehradun, viii+ 110 p.
- Kalle, R., Ramesh, T., Qureshi, Q. and Sankar, K. 2011. Density of tiger and leopard in a tropical deciduous forest of Mudumalai Tiger Reserve, southern India, as estimated using photographic capture–recapture sampling. *Acta Theriologica*, 56, 335–342.
- Kalle, R., Ramesh, T., Qureshi, Q. and Sankar, K. 2013. Predicting the distribution pattern of small carnivores in response to environmental factors in the Western Ghats. *PLoS One*, 8(11), e79295.
- Kalle, R., Ramesh, T., Qureshi, Q. and Sankar, K. 2014. Estimating seasonal abundance and habitat use of small carnivores in the Western Ghats using an occupancy approach. *Journal of Tropical Ecology*, 30, 469-480.
- Kanagavel, A., Parvathy, S., Nameer, P.O. and Raghavan, R. 2016. Conservation implications of wildlife utilization by indigenous communities in the southern Western Ghats of India. *Journal of Asia-Pacific Biodiversity*, 9(3), 271-279.
- Karanth, K. K., Gopalaswamy, A. M., Prasad, P. K., & Dasgupta, S. 2013. Patterns of human–wildlife conflicts and compensation: Insights from Western Ghats Protected Areas. *Biological Conservation*, 166, 175-185.
- Karanth, K.K., Nichols, J.D., Karanth, K.U., Hines, J.E. and Christensen, N.L. 2010. The shrinking ark: patterns of large mammal extinctions in India. *Proceedings of the Royal Society B: Biological Sciences*, 277, 1971–1979.
- Karanth, K.U. and Sunquist, M. 2000. Behavioural correlates of predation by tiger (*Panthera tigris*) and leopard (*Panthera pardus*) in Nagarhole, India. *The Zoological Society of London*, 4, 255–265.
- Karanth, K.U. and Sunquist, M.E. 1995. Prey selection by tiger, leopard and dhole in tropical forests. *Journal of Animal Ecology*, 64, 439-450.
- Karanth, K.U. and Sunquist, M.E. 2000. Behavioral correlates of predation by tiger (*Panthera tigris*), leopard (*Panthera pardus*) and dhole (*Cuon alpinus*) in Nagarhole, India. *Journal of Zoology*, 250, 255-265.
- Karthik Krishnan, M. G. (2020). Seasonality of human wildlife conflicts in Wayanad Kerala (Doctoral dissertation, Academy of Climate Change Education and Research, Vellanikkara).
- Kitchener, A. 1991. Natural history of wild cats. Comstock Publishing Associates, Ithaca, New York, xxi+280 p.
- Kolipakam, V., Singh, S., Pant, B., Qureshi, Q. and Jhala, Y.V. 2019. Genetic structure of tigers (*Panthera tigris tigris*) in India and its implications for conservation. *Global Ecology and Conservation*, 20, e00710.
- Kshetry, A. 2023 Big Cat of the Estate: Leopards in Shared Spaces, Roundglass Sustain <https://roundglassustain.com/conservation/leopards-northern-west-bengal>
- Kshetry, A., Vaidyanathan, S. and Athreya, V. 2018. Diet selection of leopards (*Panthera pardus*) in a human-use landscape in north-eastern India. *Tropical Conservation Science*, 11, 1-9.
- Kshetry, A., Vaidyanathan, S., & Athreya, V. 2017. Leopard in a tea-cup: A study of leopard habitat-use and human-leopard interactions in north-eastern India. *PLoS One*, 12(5), e0177013.
- Kshetry, A., Vaidyanathan, S., & Athreya, V. 2018. Diet selection of leopards (*Panthera pardus*) in a human-use landscape in North-Eastern India. *Tropical Conservation Science*, 11, 1940082918764635.
- Kshetry, A., Vaidyanathan, S., Sukumar, R., & Athreya, V. 2020. Looking beyond Protected Areas: Identifying conservation compatible landscapes in agro-forest mosaics in north-eastern India. *Global ecology and conservation*, 22, e00905.

- Kulkarni, D.S., 2023. Human-leopard conflict in Maharashtra | Surgical strike? India Today.
- Kumar, U., Awasthi, N., Qureshi, Q. and Jhala, Y. 2019. Do conservation strategies that increase tiger populations have consequences for other wild carnivores like leopards? Scientific Reports, 9, 14673.
- Kumara, H.N. and Singh, M. 2005. The occurrence of Rusty-spotted Cat *Prinallurus rubiginosa* (Geoffroy) in Nugu Wildlife Sanctuary, Karnataka. Journal of the Bombay Natural History Society, 102(3), 336–337.
- Kumara, H.N. and Singh, M. 2007. Small carnivores of Karnataka: Distribution and sight records. Journal of the Bombay Natural History Society, 104, 155– 162.
- Kumara, H.N., Ananda Kumar, M., Sharma, A.K., Sushma, H.S., Singh, M. and Singh, M. 2004. Diversity and management of wild mammals in tea gardens in the rainforest regions of the Western Ghats, India: A case study from a tea estate in the Anaimalai Hills. Current Science, 87(9), 1282-1287.
- Kumaraguru, A. 2006. Studies on the influence of vegetation types and prey composition on the feeding ecology of tiger, leopard and wild dog in Indira Gandhi Wildlife Sanctuary and National Park, Tamil Nadu, India. Ph.D. thesis, Bharathidasan University, Tiruchirappalli.
- Majumder, A., Sankar, K., Qureshi, Q. and Basu, S., 2013. Predation ecology of large sympatric carnivores as influenced by available wild ungulate prey in a tropical deciduous forest of Central India. Journal of Tropical Ecology, 29(5), pp.417-426.
- Majgaonkar, I., Vaidyanathan, S., Srivatsa, A., Shivakumar, S., Limaye, S. and Athreya, V. 2019. Land-sharing potential of large carnivores in human-modified landscapes of western India. Conservation Science and Practice, 2019, 1:e34.
- Mani, M. S. 1974. Ecology and Biogeography in India (M. S. Mani, Ed.; 1st ed.). Springer Dordrecht. <https://doi.org/10.1007/978-94-010-2331-3>
- Manjunatha, M. C., Swamy, I. C., & Prabhavathi, M. C. 2023. The Rising Effects of Human-leopard Conflicts in Karnataka: A Review from Media Reports. Journal of Global Ecology and Environment, 16-29.
- Marker, Laurie., & Sivamani, S. 2009. Policy for human-leopard conflict management in India. CAT news, 50(1), 23-26.
- Maroju, P.A., Yadav, S., Kolipakam, V., Singh, S., Qureshi, Q. and Jhala, Y. 2016. Schrodinger's scat: a critical review of the currently available tiger (*Panthera tigris*) and leopard (*Panthera pardus*) specific primers in India, and a novel leopard specific primer. BMC Genetics, 17(1), 1-6.
- Menon, V. 2014. Indian mammals: a field guide. Hachette Book Publishing India Pvt. Ltd., Gurgaon, India, 528 p.
- Mondal, K. 2011. Ecology of leopard (*Panthera pardus*) in Sariska Tiger Reserve, Rajasthan. Ph.D. thesis, Saurashtra University, India, 200 p.
- Mondal, K., Gupta, S., Qureshi, Q. and Sankar, K. 2011. Prey selection and food habits of leopard (*Panthera pardus fusca*) in Sariska Tiger Reserve, Rajasthan, India. Mammalia, 75, 201–205.
- Mondal, K., Sankar, K., Qureshi, Q., Gupta, S. and Chourasia, P. 2012. Estimation of population and survivorship of leopard through photographic capture-recapture sampling in west India. World Journal of Zoology, 7, 30–39.
- Mondol, S., Sridhar, V., Yadav, P., Gubbi, S., Ramakrishnan, U., 2015. Tracing the geographic origin of traded leopard body parts in the Indian subcontinent with DNA-based assignment tests. Conserv. Biol. 29, 556–564. <https://doi.org/10.1111/cobi.12393>
- Mukherjee, T., Sharma, L. K., Thakur, M., Saha, G. K., & Chandra, K. (2019). Changing landscape configuration demands ecological planning: Retrospect and prospect for megaherbivores of North Bengal. PloS one, 14(12), e0225398.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., Fonesca, G.A.B. and Kents. J. 2000. Biodiversity hotspots for conservation priorities. Nature, 403, 853-858.
- Naha, D., Chaudhary, P., Sonker, G., & Sathyakumar, S. 2020. Effectiveness of non-lethal predator deterrents to reduce livestock losses to leopard attacks within a multiple-use landscape of the Himalayan region. PeerJ, 8, e9544.
- Naha, D., Dash, S. K., Chettri, A., Chaudhary, P., Sonker, G., Heurich, M., & Sathyakumar, S. 2020. Landscape predictors of human-leopard conflicts within multi-use areas of the Himalayan region. Scientific reports, 10(1), 11129.
- Naha, D., Dash, S. K., Kupferman, C., Beasley, J. C., & Sathyakumar, S. 2021. Movement behavior of a solitary large carnivore within a hotspot of human-wildlife conflicts in India. Scientific Reports, 11(1), 3862.
- Naha, D., Sathyakumar, S., & Rawat, G. S. 2018. Understanding drivers of human-leopard conflicts in the Indian Himalayan region: spatio-temporal patterns of conflicts and perception of local communities towards conserving large carnivores. Plos one, 13(10), e0204528.
- Nath, A., Lahkar, B. P., Brahma, N., Sarmah, P., Das, A. K., Das, S., & Swargiary, A. 2021. Community, conflict and conservation: response of mammalian fauna to ecological and anthropological correlates—a critical habitat in Indo-Bhutan transboundary landscape urges multiagency cooperation.
- Nath, A., Lahkar, B. P., Brahma, N., Sarmah, P., Das, A. K., Das, S., & Swargowari, A. 2023. Breaking Dawn: Factors influencing mammalian habitat usage in western Assam following socio-political instability. Journal for Nature Conservation, 72, 126357.
- Navya, R., Athreya, V., Mudappa, D. and Shankar Raman, T.R. 2014. Assessing leopard occurrence in the plantation landscape of Valparai, Anaimalai Hills. Current Science, 107(9), 1381-1385.
- Northeast Now (2019) Assam: Leopard enters residential area in Dibrugarh, injures one <https://nenow.in/north-east-news/assam/assam-leopard-enters-residential-area-in-dibrugarh-injures-one.html>
- Northeast Now 2021 Assam: Leopard found dead in Golaghat <https://nenow.in/north-east-news/assam/assam-leopard-found-dead-in-golaghat.html>
- Northeast Now 2022 Assam: Leopard injures two in Dibrugarh <https://nenow.in/north-east-news/assam/assam-leopard-injures-two-in-dibrugarh.html>
- Northeast Now 2022 Assam: Leopard killed by villagers in Golaghat <https://nenow.in/north-east-news/assam/assam-leopard-killed-by-villagers-in-golaghat.html>

- Northeast Now 2023 Assam: Leopard killed after being hit by car in Nagaon <https://nenow.in/north-east-news/assam/assam-leopard-killed-after-being-hit-by-car-in-nagaon.html>
- Nowell, K. and Jackson, P. 1996. Wild Cats. Status Survey and Conservation Action Plan. IUCN/SSC Cat Specialist Group. IUCN, Gland, 382 p
- Patel, Shiv Kumari, Sourabh Ruhela, Suvankar Biswas, Supriya Bhatt, Bivash Pandav, and Samrat Mondol 2023. “The cost of sympatry: spatio-temporal patterns in leopard dietary and physiological responses to tiger competition gradient in Rajaji Tiger Reserve, Uttarakhand, India.” *Conservation Physiology* 11, no. 1
- Pekel, J. F., Cottam, A., Gorelick, N., & Belward, A. S. (2016). High-resolution mapping of global surface water and its long-term changes. *Nature*, 540(7633), 418-422.
- Prater, S.H. 1980. The book of Indian animals (3rd edition). Bombay Natural History Society, Bombay, xxii+324 p.
- Qamar Qureshi, Yadvendradev V. Jhala, Satya P. Yadav and Amit Mallick (eds) 2023. Status of tigers, co-predators and prey in India, 2022. National Tiger Conservation Authority, Government of India, New Delhi, and Wildlife Institute of India, Dehradun ISBN No: 81-85496-92-7
- Qureshi, Q., Gopal, R., Kyatham, S., Basu, S., Mitra, A. and Jhala, Y.V. 2006. Evaluating tiger habitat at the tehsil level. Project Tiger Directorate, Govt. of India, New Delhi and the Wildlife Institute of India, Dehradun. Technical Report TR No. 06/001, 162 p.
- Qureshi, Q., Jhala, Y. V., Yadav, S. P. and Mallick A (eds) 2023. Status of tigers, co-predators and prey in India, 2022. National Tiger Conservation Authority, Government of India, New Delhi, and Wildlife Institute of India. Dehradun. ISBN No. 81-85496-92-7
- Qureshi, Q., Saini, S., Basu, P., Gopal, R., Raza, R. and Jhala, Y.V. 2014. Connecting tiger populations for long-term conservation. Technical Report, National Tiger Conservation Authority and Wildlife Institute of India, Dehradun, TR 2014-02.
- R Development Core Team 2023. R: A language and environment for statistical computing. Computer programme. Retrieved from <http://www.R-project.org/>.
- Rahalkar, K. 2008. Attitudes of local people to conflict with leopards (*Panthera pardus*) in an agricultural landscape in Maharashtra, India.
- Ramesh, T., Kalle, R., Sankar, K. and Qureshi, Q. 2012. Spatio-temporal partitioning among large carnivores in relation to major prey species in Western Ghats. *Journal of Zoology*, 287, 269-275.
- Ramesh, T., Snehalatha, V., Sankar, K. and Qureshi, Q. 2009. Food habits and prey selection of tiger and leopard in Mudumalai Tiger Reserve, Tamil Nadu, India. *Journal of Scientific Transactions in Environment and Technovation*, 2, 170-181.
- Ramesh, T., Sridharan, N., Sankar, K., Qureshi, Q., Selvan, K.M., Gokulakannan, N., Francis, P., Narasimmarajan, K., Jhala, Y.V. and Gopal, R. 2012. Status of large carnivores and their prey in tropical rainforests of South-western Ghats, India. *Tropical Ecology*, 53(2), 137-148.
- Rana AK, Kumar N. 2023. Current wildlife crime (Indian scenario): major challenges and prevention approaches. *Biodivers Conserv.* 2023;32(5):1473-1491. doi: 10.1007/s10531-023-02577-z. Epub 2023 Mar 20. PMID: 37063172; PMCID: PMC10025790.
- Ranjitsinh, M.K. 1997. Beyond the tiger: portraits of Asian wildlife. Birajbasi Printers, New Delhi.
- Raza, R.H., Chauhan, D.S., Pasha, M.K.S. and Sinha, S. 2012. Illuminating the blind spot: a study on illegal trade in leopard parts in India (2001–2010). TRAFFIC India/WWF India, New Delhi.
- Reddy A, Vanamamalai A, Gupta S, Karanth K. 2019. Human-Wildlife conflict in Karnataka, The Need for a Human-Wildlife conflict Compensation Law. Vidhi Centre for Legal Policy, Centre for Wildlife Studies;1-45.
- Rodgers, W.A. and Panwar, H.S. 1988. Planning a wildlife protected area network in India. II Volume. Project FO: IND/82/003. FAO, Dehradun.
- Rodgers, W.A., Panwar, H.S. and Mathur, V.B. 2002. Wildlife Protected Areas in India: A Review (Executive Summary). Wildlife Institute of India, Dehradun, 44 p.
- Rodriguez, E., Morris, C.S., Belz, J., Chapin, E., Martin, J., Daffer, W. and Hensley, S. 2005. An assessment of the SRTM topographic products. Technical Report JPL D-31639, Jet Propulsion Laboratory, Pasadena, California, USA.
- Shivakumar, S., Carricondo-Sánchez, D., Athreya, V., Odden, M., Dhiman, S.P., Vaidyanathan, S., Karanth, K.K., 2023. Examining leopard attacks: spatio-temporal clustering of human injuries and deaths in Western Himalayas, India. *Front. Conserv. Sci.* 4, 1157067. <https://doi.org/10.3389/fcosc.2023.1157067>
- Sidhu, S., Raghunathan, G., Mudappa, D., & Raman, T. S. 2017. Conflict to coexistence: human–leopard interactions in a plantation landscape in Anamalai Hills, India. *Conservation and Society*, 15(4), 474-482.
- Sidhu, S., Shankar Raman, T.R. and Mudappa, D. 2015. Prey abundance and leopard diet in a plantation and rainforest landscape, Anamalai Hills, Western Ghats. *Current Science*, 109(2), 323-330.
- Singh, D. 2023. Modelling habitat suitability for the leopard in southern India through Ensemble Species Distribution Modelling (Master’s thesis, University of Twente).
- Srivastha, A., Majgaonkar, I., Sharma, S., Singh, P., Punjabi, G.A., Chawla, M.M. and Banerjee, A. 2020. Opportunities for prioritizing and expanding conservation enterprise in India using a guild of carnivores as flagships. *Environment Research Letters*, 15, 064009.
- Srivastha, A., Ramachandran, V., Saravanan, P., Sureshbabu, A., Ganguly, D. and Ramakrishnan, U., 2023. Topcats and underdogs: intraguild interactions among three apex carnivores across Asia’s forestscapes. *Biological Reviews*, 98(6), pp.2114-2135.
- Sunquist, M., and F. Sunquist. 2002. Wild Cats of the World. University of Chicago Press, Chicago and London, 452 pp, ISBN 0-226-77999-8
- The Arunachal Times 2022 Stray leopard caged; to be shifted to Itanagar Zoo. <https://arunachaltimes.in/index.php/2022/09/07/stray-leopard-caged-to-be-shifted-to-itanagar-zoo/>



- The Federal 2020 Leopard count rises, but man-animal conflict remains a worry in TN. <https://thefederal.com/news/leopard-count-rises-but-man-animal-conflict-remains-a-worry-in-tn/?infinite-scroll=1/>
- The New Indian Express 2024: Growing attacks in urban pockets of Uttarakhand cause concern. (<https://www.newindianexpress.com/nation/2024/Jan/18/growing-leopard-attacks-in-urban-pockets-of-ukhand-cause-concern>)
- The Telegraph (2021) Wild leopard killed in road accident on Asian Highway 48 in Alipurduar <https://www.telegraphindia.com/west-bengal/wild-leopard-killed-in-road-accident-on-asian-highway-48-in-alipurduar/cid/1991348>
- The Times of India 2021 Canacona's man-animal conflict saga continues. <https://timesofindia.indiatimes.com/city/goa/canacona-man-animal-conflict-saga-continues/articleshow/81519691.cms>
- The Times of India 2024. Leopard scare in Dehradun colonies after attack on 12-year-old boy. (<https://timesofindia.indiatimes.com/city/dehradun/leopard-scare-in-doon-colonies-12-year-old-boy-attacked/articleshow/106884998.cms>)
- Times of India 2023. Armed forest guards in Assam school to keep leopard at bay <https://timesofindia.indiatimes.com/city/guwahati/armed-forest-guards-in-assam-school-to-keep-leopard-at-bay/articleshow/105276721.cms>
- Times of India 2023a. Leopard strays into busy Guwahati locality, injures three <https://timesofindia.indiatimes.com/city/guwahati/leopard-strays-into-busy-guwahati-locality-injures-three/articleshow/106138473.cms>
- Uphyrkina O., Johnson W.E., Quigley H., Miquelle D., Marker L., et al. 2001. Phylogenetics, genome diversity and origin of modern leopard, *Panthera pardus*. *Molecular Ecology* 10: 2617–2633.
- Uttarakhand Forest Department 2024. Human Wildlife Conflict mitigation cell, Dehradun, Uttarakhand
- Vyas, P., & Sengupta, K. 2014. Human-leopard conflict in North Bengal, India. *Tigerpaper: Regional Quarterly Bulletin on Wildlife and National Parks Management*, 41(1), 1-6.
- Working Group on Addressing Issues Related to Human-Wildlife Interactions in Kerala. 2022. Fourteenth Five Year Plan (2022-2027). Government of Kerala, Kerala State Planning Board, Agriculture Division. <https://spb.kerala.gov.in/sites/default/files/inline-files/Human%20wildlife%20conflict.pdf>

*\*Conflict information has been collated from newspaper articles.*



©Subharanjan Sen

# Acknowledgements

We acknowledge the funding support of National Authority, CAMPA, MoEFCC. We are also thankful to Sh. Subhash Chandra, CEO, National Authority for his support. Dr. S. Sathyakumar (Registrar), Dr. Bitapi Sinha (Research Coordinator), Dr. V. P. Uniyal and Dr. S. K. Gupta (Nodal Officer Externally Aided Projects), Dr. K. Ramesh and Dr. Gautam Talukdar (Nodal Officer IT), Baljeet Kaur, Rajneesh Sharma and Narendra Singh Chauhan (Finance Officer) are acknowledged for their support and facilitation. Dr. Bilal Habib is acknowledged for providing data. Dr. Sutirtha Dutta, Dr. Amit Kumar, Dr. Navendu Page, Dr. I.P Bopanna (WWF India), and Dr. Manjari Roy are acknowledged for providing training to research biologists.

## Annexure I

### A. National Tiger Conservation Authority (NTCA) Team

**Member Secretary, NTCA:** Dr. S.P. Yadav

**Nodal Officers:** Dr. Amit Mallick, Inspector General of Forest  
Sh. Rajendra Garawad, Deputy Inspector General of Forest

#### Head Quarter, New Delhi

1. Dr. Vaibhav C. Mathur, Deputy Inspector General of Forest
2. Ms. Banumathi, G., Assistant Inspector General of Forest
3. Md. Sajid Sultan, Assistant Inspector General of Forest
4. Shri Hemant Singh, Assistant Inspector General of Forest

#### Regional Office, Bengaluru

1. Shri. N.S. Murali, Inspector General of Forest
2. Ms. Harini Venugopal, Assistant Inspector General of Forest

#### Regional Office, Guwahati

1. Shri. W. Longvah, OSD & Inspector General of Forest (retd.)
2. Ms. Agatha Momin, Biologist

#### Regional Office, Nagpur

1. Shri. Hemant Bhaskar Kamdi, Assistant Inspector General of Forest
2. Shri. Anil Dashare, Biologist

### B. Wildlife Institute of India team

**Director:** Shri Virendra R. Tiwari,

**Dean:** Dr. Ruchi Badola,

**Project Investigators:** Dr. Yadvendradev V. Jhala, Prof. Qamar Qureshi and Dr. Vishnupriya Kolipakam

**NTCA-WII Tiger Cell Scientists:** Dr. Kausik Banerjee, Dr. Ujjwal Kumar, Dr. Swati Saini, Dr. Shikha Bisht, Dr. Ayan Sadhu

**Collaborators:** Dr. Lex Hiby (Conservation Research Ltd., UK) and Dr. Saket Anand (Indraprastha Institute of Information Technology, New Delhi)

**M-STRIPEs Team:** Shri Anup Pradhan, Shri Ashish Prasad, Shri Deb Ranjan Laha, Ms. Krishna Mishra, Ms. Kainat Latafat, Shri Dhruv Jain, Shri Manish Ashok Singanjude, Ms. Rutu J. Prajapati, Shri Nanka Lakra, Ms. Amal Fathima, Shri Yash Dabholkar, Ms. Deeksha N., Ms. Sagarika Das, Shri Keshab Gogoi

**Senior Research Biologists:** Dr. Bhim Singh, Ms. Genie Murao, Dr. Anindita Bidisha Chatterjee, Ms. Shravana Goswami, Shri Jayanta Kumar Bora.

**Data Archiver and Management:** Shri G. Muthu Veerapan and Shri Harsh Deep Sethi

# Research biologists

Shri Abhilash Nair	Ms Mahajan Rashmi Deepak
Shri Abhishek Shukla	Ms Mouli Bose
Shri Aman Deep Rathi	Shri Moulik Sarkar
Shri Ananya Ajay	Ms Mridula
Shri Ananya Dutta	Ms Nupur Rautela
Ms. Ananya Sengupta	Shri Omkar Nar
Ms. Anshuman Gogoi	Ms Parul Sen
Shri Arif Ahmad	Ms Pooja Latwal
Ms Aritra Roy	Shri Prayas Auddy
Shri Ayan Khanra	Shri Rajrajeshwar Thakar
Ms Deepali Chatrath	Shri Richard S Sangma
Shri Devvrat Singh	Ms Ritu Bisht
Ms Drashti Gosai	Shri Rohan Desai
Ms Farah Naz	Ms Satemmenla Longchar
Shri Gaurav Anil Shinde	Ms Sayali Aspat
Ms Gausiya Kelawala	Shri Shahzada Iqbal
Ms Gayatri Bakhale	Shri Shivam Tiwari
Ms Geetanjali Vashistha	Ms Shrishti Joshi
Ms Harshini Jhala	Ms Shriya Milind Auradkar
Shri Himanshu Kumar	Shri Sourav Das
Ms Juri Roy	Ms Stuti Anjaria
Ms Kalpana Roy	Shri Sultan
Ms Kamakshi Singh Tanwar	Ms Sumandrita Banerjee
Shri Kathan Bandyopadhyay	Ms Susmita Nilesh Patil
Shri Kaushik Mohan Koli	Ms Udit Garbyal
Ms Kesha Patel	Ms Umang Kaur Josan
Ms Maitreyee Vishwas Bhawe	Ms Vaishnavi Gusain
Shri Mayur Markhad	Ms Vedanshi Maheshwari
Ms Meghavi Purohit	Shri Vishnuvardhan
Shri Mohit Kumar Patra	Ms Yashi Singh

# Technical Assistants

Shri Abhinav K.S. Yadav	Ms Riya Adak
Shri Abhishek Petwal	Shri Nitesh Goswami
Shri Akash Rana	Shri Pratik Majumder
Shri Ankan Halder	Shri Pratik Pramod Pansare
Shri Anurag Nashirabadkar	Ms Preeti Parihar
Ms Archana Negi	Ms Richa Joshi
Ms Arti Adhikari	Ms Ritupriya Debnath
Ms Arukshata Thakur	Shri Rohit Kumar
Shri Ashish Joseph	Shri Rudrajyoti Barman
Ms Ashwini Mamgain	Shri Rushikesh Kadam
Shri C. Jebin Bristo	Ms Saana Benurwar
Ms Chaiti Tripura	Ms Shankhamala Ghosh
Shri Dixit Verma	Ms Sheela Kanswal
Shri Gorati Arun Kumar	Shri Shiladitya Acharjee
Shri Hritik Dhami	Ms Shrawani Wankhade
Shri K. M. Sooraj Murali	Ms Shreya Yadav
Ms Kiran Rawat	Shri Shubham Kumar Maletha
Shri Lakshman Bhजारंग	Ms Shweta Singh
Shri Lalitkumar Patil	Ms Sneha Madhwal
Shri Manas Shukla	Ms Sonali Aswal
Ms Maneesha Bhatt	Ms Sonika Phogat
Ms Monal Rajendra Jadhav	Ms Subhalaxmi Muduli
Shri Mukesh Kumar	Shri Suraj Chauhan
Ms Nandita Barman	Ms Suranjita Roy
Ms Neelam Negi	Ms Sushree Subhangi Sahu
Ms Neetu Bathla	Ms Susmita Panggam
Ms Neha Kumari	Shri Swadhin Kumar Jena
Ms Neha Tolia	Ms Swati Chandola
Ms Neharika Viridi	Ms Swati Gairola
Ms Nivedita Basu	Ms Swati Kukreti
Shri Nishant Saraswat	Ms Swati Singh

Shri Nishi Nath Halder	Shri Tryambak Dasgupta
Ms Pallabi Majhi	Ms Upasna Thakur
Shri Piyush Tripathi	Ms Vartika Negi
Ms Pooja Sharma	Ms Yamini Nayal
Ms Prateeksha Nath	Shri Yashaswi Garg
Ms Rashi Nautiyal	Ms Yati Gairola

## Interns and volunteers

Shri Abhay Thakur	Ms Abhilasha Srivastava
Shri Akshay Jain	Shri Pankaj Ojha
Ms Bhawana Pant	Shri Parth Kathad
Ms Farha Usmani	Ms Pinky Yadav
Ms Geeta	Shri Pranav Thapa
Shri Harshal Waghmare	Ms Preeti Tripathi
Shri Himanshu Matta	Ms Rajeshwari Jadhav
Shri Inderjeet Singh	Ms Riddhi Sondagar
Ms Kavita Bisht	Shri Roheel Taunk
Ms Madhu Panwar	Shri Saurav Panday
Ms Mahima Gagar	Shri Shaikh Obair Aqueel Ahmad
Ms Monibhadra Roy	Shri Shantanu Ishwar Nagpure
Ms Moupika Gosh	Shri Shantanu Sharma
Ms Nivedita Sharma	Ms Sneha Mane
Ms Nivedita Singh	Shri Yajurved Barua
Ms Sasi Murugesan	

**Design:** Studio Lophius

**Extract Compare Team:** Ms. Monika Saraswat, Ms. Genie Muraio, Ms. Rutu J. Prajapati, Ms. Juri Roy, Shri. Nishant Saraswat, Ms. Upasana Thakur, Shri. Mukesh Kumar, Ms. Deepali Chatrath, Ms. Aritra Roy, Ms. Chaiti Tripura, Shri. Shivam Tiwari

**Finance:** Shri Surendra Singh Yadav

**Scientific Administrative Assistant:** Shri Harsh Deep Sethi, Ms. Pallavi Sharma

**Office Support Staff:** Shri Devendra Singh, Shri Shambhu Kimothi

## Program developers

Shri Prabir De (SCIENCE, Dehradun) and	Kaiinos Geospatial Technology Pvt. Ltd., Hyderabad
Dr. Saket Anand	Indraprastha Institute of Information Technology, New Delhi

## Chief Wildlife Wardens in the states involved in tiger and leopard status estimation 2022

S. No.	States	Name
1	Andhra Pradesh	Sh. N. Prateep Kumar, Sh. Y. Madhusudhan Reddy
2	Arunachal Pradesh	Sh. N. Tam
3	Assam	Sh. M.K. Yadava
4	Bihar	Sh. Prabhat Gupta
5	Chhattisgarh	Sh. Sudhir Agarwal, Sh. P.V. Narsingh Rao, Sh. R.K. Singh
6	Goa	Sh. Santosh Kumar, Sh. Saurabh Kumar
7	Jharkhand	Sh. P.K. Verma, Sh. Rajiv Ranjan, Sh. Ashish Rawat, Sh. Shashikar Samanta
8	Karnataka	Sh. Sanjay Mohan, Sh. Vijaykumar Gogi, Sh. Rajiv Ranjan
9	Kerala	Sh. Surendra Kumar, Sh. Devendra Kumar Verma, Sh. Bennichan Thomas, Sh. Ganga Singh
10	Madhya Pradesh	Sh. Alok Kumar, Sh. J.S. Chouhan
11	Maharashtra	Sh Mahip Gupta, Sh. Sunil Limaye
12	Mizoram	Sh. PI Lalrammawii Sailo
13	Nagaland	Sh. Ved Pal, Sh. Satya Prakash Tripathi
14	Odisha	Sh. S. K. Popli, Sh. Shashi Paul
15	Rajasthan	Sh. Arindam Tomar, Sh. M.L Meena
16	Tamil Nadu	Sh. Syed Muzammil Abbas, Sh. Shekhar Kumar Niraj, Sh. Srinivas R Reddy
17	Telangana	Smt. R Sobha, Sh. Swargam Srinivas, Sh. Rakesh Mohan Dobriyal
18	Uttar Pradesh	Sh. S. Singh, Sh. K.P. Dubey
19	Uttarakhand	Sh. J.S. Suhag, Dr. P.M. Dhakate, Dr. Samir Sinha
20	West Bengal	Sh. V.K. Yadav, Sh. Debal Ray

## All India tiger and leopard estimation Nodal Officials in the States

S. No.	States	Name
1	Andhra Pradesh	Sh. D.A. Kiran, Smt. T. Nagamaneswari
2	Arunachal Pradesh	Sh. Millo Tasser
3	Assam	Dr. Satyendra Singh
4	Bihar	Sh. Surender Singh
5	Chhattisgarh	Sh. T. Aashish
6	Goa	Sh. Jabestin A.
7	Jharkhand	Sh. Kumar Ashutosh
8	Karnataka	Sh. Subhash Malkhede, Sh. Kumar Pushkar, Sh. Biswajit Mishra
9	Kerala	Sh. P.P. Pramod, Sh. K. Vijayananthan, Sh. K.V Uthaman, Sh. P Muhammed Shabab
10	Madhya Pradesh	Dr. H.S. Negi, Sh. Subharanjan Sen, Sh. Rajnish Kumar Singh
11	Maharashtra	Sh. B.S. Hooda
12	Mizoram	Sh. Pu Laltnanhlua Zathang
13	Odisha	Sh. J.D. Pati
14	Rajasthan	Sh. Sharda Pratap Singh, Sh. Nandlal Prajapat
15	Tamil Nadu	Smt. C.H. Padma, Sh. Akash Deep Baruah
16	Telangana	Dr. D. Samhita, Sh. B. Srinivas, Sh. C.P. Vinod Kumar
17	Uttar Pradesh	Sh. Sanjay Kumar
18	Uttarakhand	Sh. Ranjan K. Mishra
19	West Bengal	Sh. Piar Chand, Sh. Rajesh Kumar

## Field directors of Tiger Reserves

State	Tiger Reserve	Name
Andhra Pradesh	Nagarjunasagar Srisailam	Sh. Y. Srinivas Reddy
Arunachal Pradesh	Pakke	Sh. Suraj Singh, Sh. Satya Prakash Singh
	Kamlang	Ms. Chesta Singh , Sh. Harsh Raj Wathore
	Namdapha	Sh. Aduk Paron, Sh. Pekom Ringu
Assam	Kaziranga	Sh. P. Shivakumar, Sh. Jitindra Sarma
	Manas	Dr. Vaibhav C. Mathur
	Nameri	Sh. Pankaj Sharma , Sh. Piraisoodan B.
	Orang	Sh. Pradipta Baruah
Bihar	Valmiki	Dr. Neshamani K., Sh. Hemkant Rai
Chhattisgarh	Achanakmar	Sh. S. Jagdisan
	Indravati	Sh. A. Srivastava, Sh. Mohammad Shahid
	Udanti Sitanadi	Sh. Rajesh Pandey, Ms. Pranita Paul, Sh. Mercy Bella

Goa	--	Sh. Jabestin A., Sh. Anand Jhadhav, Sh. Prem Kumar, Sh. Aniket Gaonkar
Jharkhand	Palamau	Kumar Asutosh
Karnataka	Bandipur	Dr. P. Ramesh Kumar
	Bhadra	Sh. Prabhakaran
	BRT Hills	Dr. Santhosh Kumar G., Ms. Deep George Contractor
	Kali	Sh. Maria Christu Raju
	Nagarhole	Sh. Mahesh Kumar, Sh. Harshakumar Chikkanaragund
Kerala	Parambikulam	Sh. K Vijayanathan, Sh. K.V. Uthaman, Sh. P. Muhammed Shabab
	Periyar	Sh. Anoop K.R., Sh. P.P. Pramod
Madhya Pradesh	Bandhavgarh	Sh. B.S. Annigeri , Sh. Rajiv Kumar Mishra
	Kanha	Sh. S.K. Singh
	Panna	Sh. Uttam Sharma, Sh. Brijendra Jha
	Pench	Sh. Deb Prasad , Sh. Ashok Kumar Mishra
	Sanjay Dubri	Sh. Y.P. Singh, Sh. Amit Dubey
	Satpura	Sh. L. Krishnamoorthy
Maharashtra	Bor	Dr. Ravikiran Govekar, Ms. Shri Laxmi
	Melghat	Ms. Jayoti Banerjee
	Nawegaon Nagzira	Sh. Manikandan Ramanujam, Sh. Jayram Gouda
	Pench	Dr. Ravikiran Govekar, Ms. Shri Laxmi
	Sahyadri	Sh. Samadhan Chauhan, Sh. N.S. Laddkat
	Tadoba Andhari	Sh. Jitendra Ramgaonkar
Mizoram	Dampa	Sh. C. Lalbiaka, Sh. Pu Zira
Odisha	Satkosia	Sh. Ashok Kumar, Sh. M. Yogajayanand
	Similipal	Sh. M. Yogajayanand, Sh. Ashok Kumar, Sh. Prakash Chand
Rajasthan	Mukundara Hills	Sh. Sedu Ram Yadav, Sh. Sharda Pratap Singh
	Ramgarh Visdhari	Sh. Sedu Ram Yadav, Sh. Sharda Pratap Singh
	Ranthambore	Sh. T.C. Verma, Sh. Sedu Ram Yadav
	Sariska	Sh. Roop Narayan Meena
Tamil Nadu	Anamalai	Sh. I. Anwardeen, Sh. S. Ramasubramanian
	Kalakad Mundanthurai	Dr. N. Senthil Kumar, Ms. R. Padmawathe
	Mudumalai	Sh. K.K. Kaushal, Sh. D. Venkatesh
	Sathyamangalam	Sh. Nihar Ranjan, Sh. S. Ramasubramanian, Sh. K. Rajkumar
	Srivilliputhur Megamalai	Sh. Deepak S. Bilgi, Dr. N. Senthil Kumar, Ms. R. Padmawathe
Telangana	Amrabad	Sh. B. Srinivas, Ms. Kshitija
	Kawal	Sh. C.P. Vinod Kumar
Uttarakhand	Corbett	Sh. Rahul, Sh. Neeraj Kumar, Sh. Dheeraj Pandey
	Rajaji	Dr. Saket Badola
Uttar Pradesh	Dudhwa	Sh. Sanjay Kumar, Sh. B. Prabhakar
	Pilibhit	Sh. Lalit Kumar Verma
	Ranipur	Sh. S.N. Mishra
West Bengal	Buxa	Sh. Buddha Raj Sewa, Sh. Apurba Sen
	Sundarban	Sh. Tapas Das, Sh. Ajoy Kumar Das



# Officials and Biologists Who Coordinated Sampling exercise in the State 2022

Sl NO	State	Name
1	Andhra Pradesh	Shri Srinivasa Reddy, Shri Alan Chong Teron, Shri Vigneesh Appavu G, Shri Vineet Kumar Nanyal, Shri Ravindra Dhama, Shri P. Venkata Sandeep Reddy, Shri C.Chaitanya Kumar Reddy, Shri Y.V.Narashimha Rao, Shri G.Satish
2	Arunachal	Shri Tapek Riba, Shri T. Pali, Shri Aduk Paron, Ms Chesta Singh, Shri Suraj Singh, Shri Harshraj Dhinkar Wathore, Shri Kime Rambia, Shri Taluk Rime, Shri Tage Mili, Shri Kenjum Membron, Shri Tajum Yomcha, Shri Chandan Ri, Shri Mayurkumar Hamubhai Variya, Shri Aditya Das, Shri Son Yowa Hade, Shri Tukum Macha, Shri Mayuk Lamgu, Shri Rubu Tado, Shri Token Rime, Shri Kelsang Dechen, Shri Deokon Chiri, Shri Julu Brah, Shri Dongro Natung, Shri Hage Tare, Shri Kame Rigom, Shri Paramod Singh Meyan, Shri Dyaba Mipi, Shri Midu Molo, Shri Dilip Molo, Shri Sagar Sonar, Shri Ema Mili, Shri Anjitey Mikhu
3	Assam	Shri P. Sivakumar, Shri Jatindra Sarma, Dr. Vaibhav C. Mathur Shri Pankaj Sharma, Shri Ramesh Kumar Gogoi, Shri Pradipta Baruah, Shri Piraisoodan B, Shri Dibakar Das, Shri Chiranjeev Jain, Shri Arun Vignesh, Shri Rabindra Sarma, Shri Pallav Kumar Deka, Shri Jayanta Deka, Shri Kanak Baishya, Shri Khagesh Pegu, Shri Chiranjeev Jain, Shri Sheshidhar Reddy, Shri Nayanj Jyoti Rajbangshi, Shri Champak Deka, Shri Khanindra Kalita Shri Nayanjyoti Rajbanshi, Shri Chakrapani Ray, Shri Rajib, Shri Jesimuddin Ahmed, Shri Bidyut Bikas Borah, Shri Bibhuthi Ranjan Gogoi, Shri Bibit Dehingia, Shri Debajit Saikia, Dr. Pranjal Gogoi, Shri Sonam Kr. Gupta, Shri Arup Kalita, Shri Jatindra Mohan Das, Shri Pradib Dev Goswami, Shri Soumitra Das, Shri Manjit Sonowal, Shri Bibhuti Majumdar, Shri Debashish Buragohain, Shri Babul Brahma, Shri Krishna Hanse, Shri Pankaj Bora, Shri Pushpadhar Burgohain, Shri Dibyajyoti Deori, Shri Nayanjyoti Borah, Shri Shahnawaj Hussain, Shri Abhinav Baruah, Shri Bastav Borkatoky, Shri Raj Pallav Neog Shri Karmeshwar Boro, Shri D Rongpi, Shri Roji Barman, Shri Basiram Brahma, Shri Rajendra Nath, , Shri Girish Kalita, Shri Deep Kalita, Shri Amrit Doley, Shri Jyotishman Deka, Shri Uddipta Kalita, Shri Sajid Choudhury, Shri Himangshu Sarma, Shri Papul Rabha, Shri Dipankar Deka, Shri Nayan Pathok, Mrs Papori Kachari Shri Bichitra Narayan Sarma, Shri Satyajit Deka,
4	Bihar	Shri Surendra Singh, Shri Hemakant Roy, Dr Nesamani K., Shri Neeraj Narayan, Shri Ambrish Kumar Mall, Shri Pradyumn Gaurav, Shri Manish Verma, Shri Amita Raj, Shri Sourabh Verma, Shri Arif Ahmad, Dr. Kamlesh K Maurya (WWF India), Shri Soham Sachin Pattekar (WWF India).
5	Chhattisgarh	Shri S.Jegadeeshan, Shri S.P.Paikra, Mrs. Praneetha Paul, Shri Ganveer Dhammshil, Shri Satyadeo Sharma, Shri Varun Jain, Mrs. Priyanka Tripathi
6	Goa	Shri Jebestin A., Sh Anand Jhadhav, Shri Prem Kumar, Shri Aniket Gaonkar
7	Jharkhand	Sri Kumar Ashutosh, Shri Kumar Ashish, Shri Prajesh kant jena, Shri Manish Kumar Bakshi, Shri Ravindra Kumar



©Achintya Singh

8	Karnataka	<p>Shri Biswajit Mishra, Shri Ramesh Kumar, P., Shri Mahesh Kumar, Shri Harshakumar Chikkanaragund., Shri Prabakaran, S., Shri Santhosh Kumar, G., Ms Deep Contractor, Shri Maria Christu Raja, D., Shri T. Heeralal, Shri S. S. Lingaraja, Shri Vijay Salimath, Shri Yatish Kumar D., Shri Manoj Kumar, Shri Upendra Pratap Singh, Shri Manjunath.R.Chavan, Shri P. Shankar, Shri S. Venkatesan, Shri B N N Murthy, Shri Prakash S. Netalkar, Dr.Malathi Priya, Dr. Hanumanthappa K T, Shri Sandip Hindurao Suryawanshi, Shri Chandrashekar Naik K, Shri N. H Jagannatha, Shri G P Harsha Bhanu, Shri Sumitkumar Subhashrao Patil, Shri Dinesh Kumar Y K, Shri Prabhakar Priyadarshi, Shri Arsalan, Shri Shivshankar .E, Shri Devaraju V, Shri Prashant Shankhinamath, Shri Prashanth, P.K.M., Dr.Ajjaiah G.R., Shri Ganapathi K, Shri Vasanth Reddy K V, Shri S. G. Hegde, Shri Gopal, Shri Nandish, L, Dr. Santhosh Kumar G. Shri N E Kranthi, Shri Nilesh Shinde, Shri Yashpal Kshirsagar, Ms Dipika Bajpai, Shri Balakrishna S, Shri Basavaraj Kn, Ms Anupama H, Ms Vaanathi M M, Ms Sareena Sikkalagar, Shri Chandranna A, Ms Kajol Ajit Patil, Shri A T Poovaiah, Shri M Sivaram Babu, Shri Sharanabasappa, Shri Ruthren P, Shri Ashish Reddy Mv, Shri Dinesh Kumar Yk, Ms Seema P A, Shri Ravishankar C, Ms Kamala, Dr V Karikalan, Shri Kapileshwara Mahadevappa Gamanagatti, Shri Mohan Kumar D, Shri G U Shankar, Shri Im Nagaraj, Shri Ramesh Kumar, P., Shri Kanteppa, Shri T. Bhaskar, Shri Kanteppa, Shri P R Manjunath, Shri Umar Badshan Kurgund, Shri S Y Bilagi, Shri Charan G S, Ashwathappa T, Shri Sahan Kumar TM, Shri Prakash A D, Shri Chandrashekar Patil, Shri Suresh Teli, Shri Mallinath Kusanal, Shri Shivarudrappa Kabadagi, Shri BV Chauvan, Shri Manjunath G Naik, Shri Sharan Kumar, Shri Kavya Chaturvedi, Shri Ashok R Bhat, Ms Himavathi Bhat, Shri SM Vali, Shri Ragu D, Shri Ragu D, Shri Ashok B Alaguru, Shri Sumith Kumar Subhashrao Patil, Shri Ravikumar, Shri Paramesh K, Shri Ankaraju M N, Shri Nagendra Prasad, Ms Bhagyalaxmi MC, Ms R Vanitha, Shri Prabhakar Priyadashi, Shri Sandeep Hindurao Suryavanshi, Shri Ramesh BR, Shri Chetan Mangala Gasti, Shri K N Manjunath, Shri K S Chengappa, Ms Ratnaprabha T A, Shri Saurabkumar, Shri B Y Yalager, Shri Suresh CN, Shri Srinivas N Yaradoni, Basvaraj Dange, Dr. Shivakumar Gajare, Shri Sunil Kumar Chauvan, Shri Md. Fayazudin, Shri Md. Asad, Shri Sathish, Shri Mahadev, Shri Gopal, DS Dayanand, Shri Moshin, Shri Nehru, Ms Roshani, Kumari Kajol Ajith Patil, Shri VP Karyappa, Shri N Subramanya Rao, Shri VP Karyappa, Shri Subramanya, Shri Rangaswamy KN, MS Anusha P, Shri BS Shashidar, Shri Shankaregouda, Shri RD Putnalli, Shri Suresh, Shri K G Prakash, Shri Shridhar R, Shri Gopya Naik, Shri CY Shivamurthy, Shri Praveen Kumar Basrur, Shri K V Subramanya, Shri Ganesh V Thadagani, Ms Sunita Mallikarjun Nimbargi, Shri Muddanna, Ms Roshni AJ, Shri Lohhith BR, Shri M. N. Naveen, Shri G. Ravindra, Shri K. Paramesh, Shri K.S. Chengappa, Ms Rathnaprabha T.A., Shri Sandeep Surya Vanshi, Shri Mahadevaiah, M., Ms Kavya Chaturvedi, Shri Shivananda S Todkar, Shri Amarakshar. V.M, Shri Mahadev S P, Shri Sathish A.V, Shri Gopal K P, Shri Amarakshar, V.M., Shri Shaikh Abdul Aleem Siddiqui, Ms Niveditha, T.P., Shri Vinay, L., Shri Imran Patel, Ms Kumari. Jeevitha, B., Ms Kumari. Varidhi Krishna.</p>
9	Kerala	<p>Shri Ganga Singh, Shri Pramod P.P., Shri Sanjayankumar, Shri R. S. Arun, Shri K. R. Anoop, Shri K. Vijayanathan, Mrs. K.S. Deepa, Shri P Muhammed Shabab, Shri Pramod P.P., Shri R. Sujith, Shri K.V Harikrishnan, Shri Abdul Assis, Shri P. K. Jayakumar Sharma, Shri Ayush Kumar Kori Shri Sunil Sahadevan, Shri A. Shanavas, Shri K. I. Pradeep Kumar, Shri Anil Antony, Shri Varun Dalia, Shri Ramesh Bishnoi Shri Vinod Kumar M G, Shri Subhash K.B, Shri N. Rajesh, Shri C.V. Rajan Shri Sambudha Majumder, Shri Ravikumar Meena Mrs. R. Lekshmi, Shri T. Aswin Kumar, Shri P. Praveen, Shri Kurra Srinivas, Shri Surjith M.K, Shri Manoj K, Shri P. Biju, Shri P. Karthick, Shri Abdul Latheef C, Mrs. A. Shajna, Shri K.J. Martin Lowel, Shri Patil Suyog Subhash Rao, Shri K. V. Harikrishnan, Shri S. V. Vinod, Shri Jayachandran G, Shri Santhosh Kumar V, Shri R. Sujith, Shri Prabhu P.M, Shri Ani J R, Shri I. S. Suresh Babu, Shri Manu Sathyan, Shri Sanil, Dr. M. Balasubramanian, Shri Vishnu Vijayan, Shri M. Ramesh Babu, Shri Anoop V, Shri Vishnu O, Shri Rahul C.M, Shri Rahul R</p>
10	Madhya Pradesh	<p>Dr. H.S. Negi, Shri Subharanjan Sen, Dr. B.S. Annigeri, Shri L. Krishnamoorthy, Shri Uttam Sharma, Sh.S. K. Singh, Shri Y. P. Singh, Shri Ashok Mishra, Shri Rajiv Kumar Mishra, Shri Brijendra Jha, Shri Amit kumar Dubey, Shri Dev A prasad J., Shri Ripudaman Bhadoriya, Shri Ravindra Mani Tripathi, Shri Lovit Bharti, Shri Adhar Gupta, Shri Rajnish Kumar Singh, Dr. Anirudh Majumder, Shri Amitabh Agnihotri, Shri Gyan Prakash Shukla, Shri Tejas Karmarkar, Shri Sumit Saha, Ms Sangeeta Kevat, Shri Jitendra Awase, Shri Uttam Singh Sastiya, Shri Amit Khanna, Shri Rajendra Singh Chauhan, Shri Madhav Uike, Shri Rajendra Singh Solanki, Shri Suresh Kusre, Shri Virendra Kumar, Shri Rajesh Ninama, Shri Ambika Prasad Maravi, Shri Shiv Kumar Kokadia, Shri Inder Singh Bare, Shri Rameshwar Uike, Shri Rameshwar Udale, Shri Upendra Dubey (WWF India), Shri Sandip Choksey (WWF India), Shri Rahul Talegaonkar (WWF India), Shri D.P. Srivastava (SFRI)</p>

11	Maharashtra	Dr. Clement Ben, Shri B. S. Hooda, Mrs. Sreelakshmi A., Dr. Ravikiran Govekar, Mrs. Sreelakshmi A., Shri Nanabhau Sitaram Ladkat, Shri Samadhan Chavan, Dr. Jitendra S. Ramgaokar, Miss. Jayoti Banerjee, Shri Jayaram Gowda, Shri R. S. Ramanujam, Shri S. Yuvraj, Shri Prabhunath Shukla, Shri Atul Deokar, Shri Kiran Patil, Shri Pramod Panchbai, Shri Thengadi, Shri Uttam Shankar Sawant, Shri Vishal Mali, Shri G. Guruprasad, Shri Kushagra Pathak, Shri Nandkishor Kale, Shri Navalkishore Reddy, Shri Sumant S. Solanke, Mrs. Divya Bharti M., Shri N Jaykumaran, Madhumitha S., Shri Manoj N. Khairnar, Shri A.W. Nimje, Shri Kulraj Singh, Miss. Punam Pate, Shri Pawan Jeph, Shri Suresh Salunke, Shri Ganesh Patole, Shri Tushar Dhamdhare, Shri Abhijit Waykos, Shri Babu Chagan Yele, Shri Mahesh Chagan Khore, Shri R.R. Kulkarni, Shri Kamlesh Patil, Shri Indrajeet n. Nikam, Shri Ravindra. B. Kondawar, Shri Shupal Pawar, Shri Sandip Kumbhar, Shri Balkrushna Hasabnis, Shri Nandkumar Nalawade, Shri Dnyaneshwar Rakshe, Shri Sandip Jopale, Sau. Miss. Swati Vijay Maheshkar, Shri Bau- rao Krishnarao Tupe, Sau. Yogita V. Atram (Madavi), Shri Ghanshyam Raghunath Naigamkar, Shri Santosh Ramdas Thipe, Shri Kiran Wasudeo Dhankute, Shri Ravindra Haridas Chowdhari, Shri Satish Kisan Shende, Miss. Shubhangi Ravindra Krishnapurkar, Shri Pradip Laxman Chawhan, Shri Rundan Sada Shriv Katkar, Shri Arunkumar Ramlakhan Gound, Shri Yogesh V. Tapas, Shri Vivek Yewatkar, Shri Abhay Chandel, Shri Shantanu Sharma, Miss. Abhilasha Shrivastav, Shri Akash Patil, Miss. Prajakta Hushangabadkar, Shri Sachin Nikesar, Shri Sahbaz Sheikh, Shri Pawan Uttamrao Tikhile, Shri Sunil Kamdi, Shri Akash Sarada, Shri Gajanan Dhadse, Shri P. E. Patil,
12	Mizoram	Shri C. Lalbiaka, Shri Lalnunzira, Shri Andrew Lalthlamuana, Shri James Thanmawia, Shri Zoliansanga
13	Nagaland	Shri T. Aochuba, Dr. Sentitula, Shri Suman Sivasankar Sivachar W. M., Shri Rajesh Kumar, Dr. Prabhat Kumar, Shri Tokaho Kinimi, Shri Ltu, Shri Limaba, Shri Sashilemla, Shri Ailong Phom, Shri Temjenmongba, Shri Shilu, Shri Moakumdang, Shri Imkongmar, Shri P. Bendangmongba, Shri Chubanunsang, Shri Nahwang, Lansothung Lotha, Shri Wopansao, Shri Bokato, Shri Aaron Yimchunger, Shri Imnawapang
14	Odisha	Shri M. Yogajayanand, Shri Aksshay Kumar Patnayaik, Shri T. Ashok Kumar, Dr. Jagyandatt Pati, Dr. Prakash Chand Gogineni, Dr. Smrat Gowda D.S., Shri Sudhanshu Sekhar Khora, Shri Sai Kiran D.N., Shri Saroj Kumar Panda, Shri Samir Kuamr Satpathy, Ms. Anshu Pragyan Das, Shri Bimal Prasanna Acharya, Bidya Sagar, Shri Pradeep Kumar Dey, Shri Nikesh Kumar Mahapatra, Shri Samresh Kumar Biswal, Shri Bhakta P. Rath, Dr. Nimai Charan Palei, Shri Harshvardhan Singh Rathore, Shri Gatikrishna Behera, Shri Pankaj Kumar Das
15	Rajasthan	Shri Sedu Ram Yadav, Shri R. N. Meena, Shri T. C. Verma, Shri R. K. Khairwa, Shri Mahendra Kumar Sharma, Shri Bijo Joy, Shri Alok Gupta, Shri Sudarshan Sharma, Shri Sangram Singh Katiyar, Shri Sanjeev Kumar Sharma, Shri D. P. Jagawat, Shri Manas Singh, Shri Arun Kumar D, Shri Maria Shine, Shri Arvind Kumar Jha, Dr. Ramanad Bhakar, Shri Vinod Rai, Shri Kishore Singh, Shri Bheru Singh Rathore, Shri kishan Singh Ranawat, Shri Devendra Kumar Purohit Shri jayanti lal Garasiya, Shri Tarun Kumar Mehra, Shri Anurag Bhatnagar, Shri Hari Mohan Meena, Md. Mairaj, Shri N. Gokulakkannan, Shri Devendra Singh, Shri Rajrajeshwar Thaker, Shri Prayas Auddy, Shri Kaushik Koli, Ms. Deepali Chatrath, Shri Mohit Kumar Patra, Ms. Stuti Anjaria, Ms. Aritra Roy, Ms. Gayatri Bakhale, Ms. Suman-drita Banerjee, Shri Abhishek Petwal, Shri Devrat Singh, Shri Pankaj Ojha, Shri Omkar Nar, Shri Rohan Desai, Shri Gaurav Shinde, Ms. Oindrila Paul, Ms. Preeti Tripathy
16	Tamil Nadu	Shri S. Ramasubramanian, Shri Thiru. D. Venkatesh, Shri Nihar Ranjan, Shri S. Ramasubramanian, Shri Thiru. Deepak S. Bilgi, Dr. N. Senthil Kumar, Shri R. Padmawathe, Shri A. S. Marimuthu, Shri S. Arokiaraj Xavier, Shri M. G. Ganeshan, Dr. P. K. Dileep, Shri S. N. Tejesvi, Shri Tmt. C. H. Padma, Shri Tmt. C. Vidhya, Shri Selvi. S. Senbagapriya, Dr. Bhosale Sachin Tukaram, Shri Thiru. L. C. S. Srikanth, Shri Thiru. P. Arunkumar, Shri R. Kirubashankar, Shri Devendra Kumar Meena, Shri Kulal Yogesh Vilas, Shri S. Anand, Dr. H. Dileep Kumar, Mrs. C. Vidhya, Dr. J. R. Samartha, Dr. Gurusamay Dubbala, Shri A. Anbu, Shri D. Rameswaran, Shri N. Sridharan, Shri Ashok Kumar, Shri M. Ilayaraja, Shri K. Ganesh Ram, Shri M. Dhayanathan, Shri L. Sivakumar, Shri N. Manojkumar, Shri N. Rajendran, Shri Kanthan, Shri N. Mariyappan, Shri S. Murali, Shri R. Ramesh, Shri C. Sivakumar, Shri K. Suresh, Shri P. Ganeshpandiyan, Shri Dr. V. Saravanan, Shri C. Sivakumar, Shri S. Sivakumar, Shri D. Dinesh, Shri S. Sathish, Shri M. Ramalingam, Shri C. Sakthivel, Shri J. Peter Prem Chakravarthi, Shri N. Sridharan, Shri Karthik Thamizharasu, Shri K. Anvar, Dr. K. Mahesh Kumar, Shri C. Sakthivel, Shri G. Karthikeyan, Shri J. Yogesh, Shri M. Palanisamy, Dr. P. Santhosh Kumar, Dr. A. Samson, Shri Thiru. M. Parthiban, Shri Thiru. D. Keerthivasan, Miss. Thiru. S. Meenakshi Sundaram, Miss. Tmt. R. Nagalakshmi, Shri Thiru. R. Muniyandi, Shri Thiru. M. Murugan, Mrs. S. Agnes Jeya Packiavathi, Shri P. Yogeswaran, Shri J. Dalson Mani, Shri Tmt. S. Buvanewari, Shri P. Gunalan, Dr. R. Kanthasamy.
17	Telangana	Shri B. Srinivas, IFS, Sri M. J. Akbar, Ms Sunitha Bhagavath, Smt N. Kshitija, P.V. Raja Rao, Shri D. Bheema, Shri C.P. Vinod Kumar, G. Ramalingam, Sri G. Kista Goud, Sri Sivala Rambabu, Sri D. Venkateshwar Reddy, Sri A. Venkateswarlu, Sri S. Satyanarayana, Sri P. Ramakrishna, Shivani Dogra, Sri Vikas Meena, Sri S. Shantharam, Sri Petla Rajashekar, Shri Jogu Yellam, Sri S. Mahaboob, Sri Narsimha, Sri M. Ravi Kanth, Sri Md. Saleem,

18	Uttarakhand	Shri Rahul, Shri Naresh Kumar, Dr. Dheeraj Pandey, Dr. Saket Badola, Shri Siddhant Umaria, Mrs Kalyani, Shri Neeraj Sharma, Shri Digant Nayak, Shri Sher Singh, Shri Bhuvan Chand Upreti, Shri Rajendra Kumar Bisht, Shri Neeraj Sharma, Shri Prakash Chandra Arya, Shri Kishan Chand, Ramakant Tiwadi, Shri Brij Bharti Sharma, Shri LR Nag, Shri Amit Goswami, Shri Harish Negi, Mrs Shalini Joshi, Shri Lalit Mohan Arya, Mrs. Prema Tiwadi, Shri Shiv Shankar Giri, Shri Neeraj Negi, Shri Ashu Saini, Shri Sanjay Pandey, Dr. I.P. Bopanna (WWF India), Dr. Meeraj Anwar (WWF India), Shri Devrishi Chattopadhyay (WWF India), Shri Devrat Panwar (WWF India), Shri Siddhant Umaria (WWF India), Shri Yash Kumar Agarwal (WWF India), Shri Ankur Patra (WWF India), Shri Arpan Gupta (WWF India), Shri Jagjivan Dhama (WWF India), Shri Puran Singh (WWF India), Shri Paramdeep Singh (WWF India), Shri Sajal Singh (WWF India), Shri Shrestha Singh (WWF India), Miss Ankita Chand (WWF India), Miss Riya Adak (WWF India).
19	Uttar Pradesh	Shri Sanjay Kumar Pathak, Shri Kailash Prakash, Shri Sameer Kumar, Shri Sundresha, Shri Akashdeep Bhadwan, Shri Naveen Khandelwal, Shri Sitanshu Pandey, Shri Satyapal Singh, Shri Girdhari Lal, Shri Sobran Lal, Shri Shivbabu Saroj, Shri Vimlesh Kumar, Shri Ram ji, Shri Shatrohan Lal, Shri Chandrabhal Singh, Shri Ramkumar, Shri Surendar Kumar, Shri Wazir Hasan, Shri Arun Mohan, Shri Pramod Kumar Srivastav, Shri Radheysyam, Shri Mobeen Arif, Shri Apoorv Gupta, Shri G P Rana, Shri Rambharan Yadav, Shri Angad Prasad Singh, Shri Mayank Pandey, Shri Mohit Singh, Shri Liladhar, Shri Radheysyam, Shri Ramphool Yadav, Shri Amit Gangwar, Shri Rakesh Kumar, Shri Chandprakash, Shri Vijay Pal, Shri Dulare, Shri Ajay Kumar, Shri Uday Pratap Singh, Shri Manoj Kumar, Shri Krishan Kumar Pal, Shri Dharmenda Dhaka, Shri Ramkant Pandey, Dr. Mudit Gupta, Shri Ashish Bista, Shri Rohit Ravi (WWF India), Shri Dabeer Hasan (WWF India), Shri Naresh Lodhi (WWF India), Shri Kandhai Lal (WWF India), Shri Anil Kumar Srivastav (WWF India), Shri Devendar Kumar (WWF India), Shri Premchand (WWF India), Shri Ramlakhan (WWF India), Shri Mukesh Kumar (WWF India), Shri Rahul Kumar (WWF India), Shri Ravi Saxena (WWF India), Shri Mohd. Shahid (WWF India).
20	West Bengal	Shri Piar Chand, Shri Rajesh Kumar, Shri Rajendra Jakher, Shri Buddha Raj Sewa, Shri Apurba Sen, Shri Harish, Shri Sheik Fareed J., Shri Umar Imam, Shri Deepak M., Shri Parveen Kaswan, Shri Dwijaa Pratim Sen, Shri Manish Jha, Shri Nabi Kanth Jha, Shri Pallab Mukherjee, Shri Ranjan Talukdar, Shri Debdarshan Roy, Shri Aqib Alam, Shri Gebu Lepcha, Shri Novojit De, Shri D. K. Jha, Shri Balam Panja, Shri Deepak Mandal, Shri Partha Mukherjee, Ms. Arpita Patra, Shri V. Chakraborty, Shri D. Roy, Shri Partha Sarathi Pramanik, Shri Amalendu Maji, Shri Arnab Das, Shri Arnab Choudhary, Shri Sujit Kr Barma, Shri Narendranath Dutta, Shri Chinmoy Barman, Shri Anup Kar, Shri Himadri Debnath, Shri Swapan Kumar Majhi, Shri Sribas Sarkar, Shri Dhiraj Kami, Shri Biswajit Bishoi, Shri Sandip Das, Shri Nur Islam, Shri Ankan Nandi, Shri Raj Kumar Saha, Shri Biplab Ghosh, Shri Subhayu Saha, Shri Ashim Kumar Dandapat, Shri Monojit Chakraborty, Shri Debasish Mondal, Shri Amitesh Satpathy, Shri Soham Mitra, Shri Subhayu Saha, Shri Prabhat Kr Barman, Shri Promit Lal, Shri Mayukh Ghose, Shri Shekhar Sarkar, Shri Orvill Nazarearth, Ms Shikha Jasrotia.

## Annexure II

Across the country, five major workshops were held, while an additional 70 workshops were conducted online. Moreover, 35 training sessions took place in various states. States have further conducted workshops in their Protected Areas and divisions.

### List of Training of Trainers Workshops

Date of Workshop	Venue of the Workshop	Participating States	Approximate number of personnel trained
August 06 - 08, 2021	Mudumalai Tiger Reserve, Tamil Nadu	Andhra Pradesh, Goa, Karnataka, Kerala, Tamil Nadu, Telangana	112
August 12 - 14, 2021	Ranthambore Tiger Reserve, Rajasthan	Chhattisgarh, Madhya Pradesh, Maharashtra, Odisha, Rajasthan	105
August 25 - 27, 2021	Rajaji Tiger Reserve, Uttarakhand	Bihar, Uttar Pradesh, Uttarakhand, Jharkhand	80
September 02 - 04, 2021	Manas Tiger Reserve, Assam	West Bengal (North), Assam, Mizoram, Meghalaya, Nagaland, Arunachal Pradesh	96
November 02 - 04, 2021	Sundarbans Tiger Reserve, West Bengal	Staff of Sundarbans of India and Bhitarkanika NP (Odisha)	62

## Annexure III

Contribution is based on work done for field sampling, data analysis supervision, and writing up; The WII team names are in no particular order; the data presented belongs to the NTCA, WII, and State Forest Departments according to the respective MoUs;

### List of contributors

#### Shivalik Hills and Gangetic Plains Landscape

##### NTCA TEAM

Satya P. Yadav, Amit Mallick, Rajendra Garawad, Hemant Singh

##### WII TEAM

Qamar Qureshi, Yadvendradev V. Jhala, Vishnupriya Kolipakam, Shikha Bisht, Ujjwal Kumar, Swati Saini, Anindita Bidisha Chatterjee, Anup Kumar Pradhan, Ashish Prasad, Sudip Banerjee, Kainat Latafat, Monika Saraswat, G.Muthu Veerappan, Vaishnavi Gusain, Dhruv Jain, Pooja Choudhary, Abhishek Shukla, Ananya Ajay, Ananya Dutta, Anshuman Gogoi, Ayan Khanra, Devvrat Singh, Gayatri Bakhale, Geetanjali, Juri Roy, Kaushik Koli, Nanka Lakra, Mouli Bose, Mridula, Nupur Rautela, Omkar Nar, Prayas Auddy, Rajrajeshwar Thakar, Richard S. Sangma, Ritu Bisht, Sagarika Das, Shivam Tiwari, Stuti Anjaria,

Vedanshi Maheshwari, Sumandrita Banerjee, Yashi Singh, Abhishek Petwal, Anurag Nashirabadkar, Archana Negi, Ashwini Mamgain, Hrithik Dhama, Gaurav Naithani, Md. Shaan, Mukesh Saraswat, Neetu Bathla, Neha Toila, Nibedita Basu, Nishant Saraswat, Rashmi Mahajan, Rohit Kumar, Srishti Joshi, Shankhmala Ghosh, Shiladitya Acharjee, Sonika Phogat, Sonali Aswal, Swati Chandola, Swati Kukreti, Swati Singh, Upasana Thakur, Geeta, Neelam Negi, Pranav Thapa, Prateeksha Nath, Harsh Deep Sethi, Suraj Chauhan, Surbhi.

## Central India & Eastern Ghats Landscape

### NTCA Team

Satya P. Yadav, Amit Mallick, Rajendra Garawad, Hemant Kamdi, Anil Kumar Dashahre

### WII Team

Qamar Qureshi, Yadvendradev V. Jhala, Vishnupriya Kolipakam, Ujjwal Kumar, Rutu J. Prajapati, Krishna Mishra, Ashish Prasad, Omkar Nar, Manish A. Singanjude, Ayan Sadhu, Ninad A. Mungi, Shravana Goswami, Jayanta Kumar Bora, Swati Saini, Bhim Singh, Kainat Latafat, Deb Ranjan Laha, Anup Kumar Pradhan, Monika Saraswat, G. Muthu Veerappan, Keshab Gogoi, Nanka Lakra, Genie Murao, Abhishek Shukla, Gaurav Anil Shinde, Juri Roy, Rudrajyoti Barman, Anshuman Gogoi, Ayan Khanra, Sumandrita Banerjee, Richard S. Sangma, Stuti Anjaria, Dhruv Jain, Vaishnavi Gusain, Aritra Roy, Deepali Chatrath, Rajrajeshwar Thakar, Ritu Bisht, Arif Ahmad, Mohd. Akram, Monal Rajendra Jadhav, Pooja Sharma, Rohan Desai, Ashish Joseph, Lakshman Bhajarang, Shweta Singh, Subhalakshmi Muduli, Yash Dabholkar, Kiran Rawat, Rushikesh Kadam, Lalitkumar Patil, Upasna Thakur, Swati Kukreti, Geeta Negi, Nishant Saraswat, Drashti Gosai, Roheel Taunk, Susmita Panggam, Parth Kathad, Susmita Nilesh Patil, Devvrat Singh, Kaushik Mohan Koli, Sneha Madhwal, Kamakshi Singh Tanwar, Amandeep Rathi, Gitanjali Vashistha, Inderjeet Singh, Shahzada Iqbal, Shantanu Ishwar Nagpure, Mahima Gagar, Sonali Aswal, Sushree Subhangi Sahu, Swadhin Kumar Jena, Swati Singh, Vedanshi Maheshwari, Mukesh Kumar, Abhay Thakur, Preeti Parihar, Hrithik Dhama, Tryambak Dasgupta, Abhishek Petwal, Gayatri Bakhale, Mohit Kumar Patra, Pankaj Ojha, Yashi Singh, Shaikh Obair Aqueel Ahmad, Prayas Auddy, Nivedita Sharma, Sheela Kanswal, Amal Fathima, Ananya Dutta, Pinky Yadav, Chaiti Tripura, Shriya Milind Auradkar, Meghavi Purohit, Shiladitya Acharjee, Neetu Bathla, Rajeshwari Jadhav, Sasi M., Mridula, Harsh Deep Sethi, Neharika Virdi, Prateeksha Nath, Swati Chandola, Parul Sen, Nupur Rautela, Moulik Sarkar, Akshay Jain, Kesha Patel, Sultan, Harshini Jhala, Pooja Chaudhary, Kathan Bandyopadhyay, Shivam Tiwari, Sagarika Das, Arukshata Thakur, Ashwini Mamgain, Akash Rana, Ananya Ajay, Archana Negi, Neha Tolia, Nibedita Basu, Shrawani Wankhade, Pratik Pansare, Shankhamala Ghosh, Richa Joshi, Aarti Adhikari, Suraj Chauhan, K. M. Sooraj Murali, Preeti Tripathi, Umang Kaur Josan, Piyush Tripathi, Abhinav K.S. Yadav.

## Western Ghats landscape

### NTCA Team

Satya P. Yadav, Amit Mallick, N S Murali, Rajendra Garawad, Harini Venugopal

### WII Team

Qamar Qureshi, Yadvendradev V. Jhala, Vishnupriya Kolipakam, Ayan Sadhu, Genie Murao, Kausik Banerjee, Prayas Auddy, Ananya Dutta, Deepali Chatrath, Kainat Latafat, Kamakshi Singh Tanwar, Dhruv Jain, Vaishnavi Gusain, Ashish Prasad, G. Muthu Veerappan, Monika Saraswat, Manish Singanjude, Nanka Lakra, Pooja Choudhary, Rajrajeshwar Thakar, Gayatri Bakhale, Aritra Roy, Kaushik Mohan Koli, Stuti Anjaria, Abhishek Petwal, Abhishek Shukla, Sumandrita Banerjee, Vedanshi Maheshwari, Pratik Pansare, Suranjita Roy, Akash Rana, Sheela Kanswal, Anurag Nashirabadkar, Rajeshwari Jadhav, Sasi M., Sonika Phogat, Omkar Nar, Ayan Khanra, Umang Kaur Josan, Piyush Tripathi, Mukesh Kumar, Gaurav Anil Shinde, Maitreyee Vishwas Bhave, Mohit Kumar Patra, Mridula, Pankaj Ojha, Shaikh Obair Aqueel Ahmad, Harshal Waghmare, Mouli Bose, Monibhadra Roy, Nivedita Singh, Neetu Bathla, Neelam Negi, Nishi Nath Halder, Shweta Singh, Tryambak Dasgupta, Manas Shukla, Anshuman Gogoi, Riddhi Sondagar, Susmita Patil, Yashi Singh, Richard Sangma, Rohan Desai, Shivam Tiwari, Juri Roy, C. Jebin Bristo, Hritik Dhama, K. M. Sooraj Murali, Manoranjan Parida, Md. Akram, Prateeksha Nath, Pratik Majumder, Shiladitya Acharjee, Subhalaxmi, Suraj Chauhan, Sushree Subhangi Sahu, Swadhin Kumar Jena, Swati Singh, Kalpana Roy, Amal Fathima, Ananya Ajay, Harshdeep Sethi, Swati Chandola, Swati Kukreti, Ms Ritu Bisht, Udit Garbyal, Upasna Thakur, Chaiti Tripura, Neha Kumari, Pinky Yadav, Moupika Gosh, Preeti Tripathi, Shankamala Ghosh, Ritupriya Debnath.


## North Eastern Hills and Brahmaputra Flood Plains Landscape

### NTCA Team

Satya P. Yadav, W. Longvah, Rajendra Garawad

## WII Team

Qamar Qureshi, Yadvendradev V. Jhala, Vishnupriya Kolipakam, Deb Ranjan Laha, Shikha Bisht, Ujjwal Kumar, Swati Saini, Ashish Prasad, G. Muthu Veerappan, Monika Saraswat, Genie Murao, Rutu Prajapati, Richard S. Sangma, Anshuman Gogoi, Vaishnavi Gusain, Dhruv Jain, Omkar Nar, Kainat Latafat, Bhim Singh, Abhishek Petwal, Rajrajeshwar Thakar, Juri Roy, Shankhamala Ghosh, Nanka Lakra, Nishant Saraswat, Vedanshi Maheshawari, Pratik Pansare, Sneha Madhwal, Archana Negi, Ritu Bisht, Susmita Panggam, Anurag Nashirabadkar, Yash Dabholkar, Ananya Pandey, Rohit Kumar, Mridula, Amal Fathima, Deeksha N., Shiladitya Acharjee, Yashi Singh, Shivam Tiwari, Farah Usmani, Gausiya Kelawala, Ananya Sengupta, Farah Naz, Vishnuvardhan, Shrishti Joshi, Pooja Latwal, Susmita Nilesh Patil, Sonika Phogat, Neha Kumari, Sagarika Das.



©Achintya Singh









 भारतीय वन्यजीव संस्थान  
Wildlife Institute of India



  
आज़ादी का  
अमृत महोत्सव

National Tiger Conservation Authority  
B-1 Wing, 7th Floor, Pt Deendayal Antyodaya  
Bhawan, CGO Complex,  
Lodhi Road, New Delhi 110 003, India  
<https://ntca.gov.in/>

Wildlife Institute of India  
Chandrabani, Dehradun - 248001  
Uttarakhand, India  
[www.wii.gov.in](http://www.wii.gov.in)